

Material	Nitrogen (N) %	Phosphoric acid (P) %	Potash (K) %
Bulky organic materials			Fotasii (K) %
Alfalfa hay	2.5	0.5	2.0
Bean straw	1.2	0.3	1.2
Crabgrass	0.66	0.19	0.71
Grain straw	0.6	0.2	1.0
Oak leaves	0.80	0.35	0.15
Peanut hulls	1.5		0.8
Peat	2.3	0.4	0.8
Sawdust	0.2		0.2
Seaweed (kelp)	0.6		1.3
Manures			
Bat guano	10.0	4.5	2.0
Cow manure, dried	1.3	0.9	0.8
Cow manure, fresh	0.5	0.2	0.5
Hen manure, dried, with litter	2.8	2.8	1.5
Hen manure, fresh	1.1	0.9	0.5
Horse manure, fresh	0.6	0.3	0.5
Pig manure, fresh	0.6	0.5	0.4
Sheep manure, dried	1.4	1.0	3.0
Sheep manure, fresh	0.9	0.5	0.8

Kansas State University Agricultural Experiment Station and Cooperative Extension Service

Material	Nitrogen (N) %	Phosphoric acid (P) %	Potash (K) %
Rock powders			
Basic slag		8-17	
Greensand (glauconite)		1.4	4.0-9.5
Hybro-tite		0.002	
Rock phosphate (apatite)		38-40	4.5
Vegetative and animal materials			
Bonemeal, steamed	2.0	22	
Castor pomace	6.0	1.9	0.5
Cocoa shell meal	2.5	1.5	2.5
Cottonseed meal	6.0	3.0	1.0
Dried blood meal	13.0	1.5	0.8
Feathers	15.0	30.0	
Fish meal	10.0	6.0	
Fish scrap	5.0	3.0	
Garbage tankage	1.5	2.0	0.7
Hoof and horn meal	12.0	2.0	
Soybean meal	7.0	1.2	1.5
Tankage, animal	9.0	6.0	
Tankage, processed	7.0	1.0	0.1
Wood ashes		1.8	5.0

Ward Upham Horticulturist



Publications from Kansas State University are available at *www.bookstore.ksre.ksu.edu*.

Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, credit the authors, *Approximate Composition of Select Natural Fertilizer Materials*, August 2017.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, John D. Floros, Director.