



Keeping
Up With
Research
27

DECEMBER 1976

Summer Injection of 2,4-D and Tordon Herbicides to Control Unwanted Trees in Kansas Woodlands¹

**Wayne A. Geyer, Research Forester
Larry E. Biles, Area Forester**

Kansas woodlands often need to be upgraded by removing unwanted, low-quality trees. Selectively removing or killing them promotes growth of the high-quality ones remaining.

Controlling undesirable trees with herbicides is an accepted forestry practice, but new chemicals must be evaluated to find the safest and most effective ones. As herbicide results vary from region to region and among species and climates, we conducted a series of tests on typical Kansas hardwood trees. Earlier results had shown that Tordon* 101 Mixture was an effective herbicide (Geyer and Biles, 1975).

The Studies and Results

We treated trees at three sites in northeastern Kansas. Two were black walnut sapling-pole stands on intermittent streams and one was a large pole-small sawtimber cottonwood stand on the Kansas River.

The herbicides killed a high percentage of trees at all locations. All were injected through the bark at 3-inch

1. Contribution no. 586-S, Department of Horticulture and Forestry, Kansas Agricultural Experiment Station, Kansas State University, Manhattan, Kansas 66506.

* Trademark of the DOW Chemical Co. Mention of trade names is for better understanding only; no endorsement of products named is intended.

AGRICULTURAL EXPERIMENT STATION
Kansas State University, Manhattan
Floyd W. Smith, Director

intervals around bases of the trees with a Jim-Gem metering injector (Figure 1). About one milliliter of herbicide was injected into each cut (Figure 2) in late spring or early summer after the trees had leafed out.



Figure 1.—Basal metering injector we used.



Figure 2.—We made cuts 3 inches apart close to the ground and metered 1 milliliter of herbicide into each cut.

We injected three herbicides: 2,4-D, Tordon 22K, and Tordon 101 Mixture. The concentrated amine form of 2,4-D at 4 pounds acid equivalent per gallon was used. Tordon 22K, a potassium salt formulation containing 2 lb. ai/gal. of picloram, was diluted with 3 parts water. Tordon 101 Mixture, a marketed combination of 2,4-D and picloram was mixed 1:1 with water. All are water soluble and generally considered more effective than ester formulations of the same chemicals injected into a tree's vascular system.

Only Tordon 101 Mixture was used in the two walnut stands; all three herbicides were used in the cottonwood stand.

Seventeen species (977 trees) were treated: American elm (*Ulmus americana* L.), black cherry (*Prunus serotina* Ehrh.), black walnut (*Juglans nigra* L.), black willow (*Salix nigra* Marsh.), box elder (*Acer negundo* L.), cottonwood (*Populus deltoides* Marsh.), dogwood (*Cornus stolonifera* Michx.), hackberry (*Celtis occidentalis* L.), hawthorn spp. (*Crataegus* L.), honey locust (*Gleditsia triacanthos* L.) Kentucky coffeetree (*Gymnocladus dioica* [L.] K. Koch), mulberry spp. (*Morus* L.), osage orange (*Maclura pomifera* [Raf.] Schneid), red bud (*Cercis canadensis* L.), red elm (*Ulmus rubra* Muhl.), silver maple (*Acer saccharinum* L.), and sycamore (*Platanus occidentalis* L.).

Crown kill was observed during the first and second growing seasons. More than 90% of the injected trees the first year, and still more the second year had 100% crown kill from Tordon 101 Mixture (Table 1). Silver

Table 1.—Effects of Tordon 101 Mixture on several hardwood species after two growing seasons.¹

| Species | Crown Kill, % | | No. of Trees Treated |
|---------------------------|--------------------------|----------|----------------------|
| | Substantial ² | Complete | |
| American elm | 3 | 94 | 159 |
| Black cherry | | 100 | 2 |
| Black willow | | 100 | 20 |
| Black walnut | 4 | 93 | 256 |
| Box elder | 12 | 88 | 8 |
| Cottonwood | | 100 | 59 |
| Dogwood | | 100 | 2 |
| Hackberry | 25 | 70 | 20 |
| Hawthorn | | 100 | 5 |
| Honeylocust | 1 | 91 | 161 |
| Kentucky coffeetree | | 100 | 11 |
| Mulberry | 6 | 94 | 16 |
| Osage orange | 19 | 78 | 32 |
| Red bud | | 100 | 5 |
| Red elm | 12 | 88 | 25 |
| Silver maple | | 64 | 11 |
| Sycamore | | 100 | 1 |
| All species, avg. | 3 | 92 | 793 |

1. Tordon 101 Mixture 1:1 with water basal-injected at 3-inch intervals during the summer at 3 sites.

2. Substantial 1/2 to nearly complete.

maple, with a 64% crown kill appears to be somewhat difficult to control. Putting injections at closer spacing than 3 inches apart might improve its effectiveness on silver maple. Only a few trees produced basal stem sprouts after two growing seasons.

Picloram (Tordon 22K), 2,4-D (Formula 40), and picloram plus 2,4-D (Tordon 101 Mixture) were compared on 4 species in the cottonwood stand (Table 2). Only 2,4-D

Table 2.—Effects of 2,4-D, Tordon 22K, and Tordon 101 Mixture after 2 growing seasons.¹

| Species | Herbicide Injected | | | | | |
|-------------------|--------------------|-----------|------------|-----------|------------|-----------|
| | 2,4-D | | Tordon 22K | | Tordon 101 | |
| | % Complete | No. Trees | % Complete | No. Trees | % Complete | No. Trees |
| Black willow ... | 100 | 12 | 100 | 32 | 100 | 20 |
| Cottonwood | 100 | 45 | 100 | 57 | 100 | 59 |
| Mulberry | 100 | 1 | 100 | 3 | 100 | 6 |
| Silver maple | 9 | 11 | 100 | 17 | 64 | 11 |
| Sycamore | | | 100 | 3 | 100 | 1 |

1. Concentrated 2,4-D amine, Tordon 22K with 3 parts water; Tordon 101 Mixture 1:1 with water, each basal-injected at 3-inch intervals during growing season.

amine alone was ineffective on silver maple, while Tordon 101 Mixture was better (64% complete crown kill). Straight picloram (Tordon 22K) was 100% effective. Black willow and cottonwood appear to be easy to kill by any of the 3 herbicide formulations.

Discussion

Apparently basal-injected, Tordon herbicides (Tordon 101 Mixture and Tordon 22K) will effectively control undesirable hardwood trees in eastern Kansas. Both are more effective than 2,4-D amine alone on the hard-to-kill silver maple during the growing season.

Comparative trials of 2,4-D amine and Tordon 101 Mixture on a variety of hardwood species in the mountains of Arkansas showed picloram effective on many species, including some resistant to 2,4-D alone (Ferguson and Lawson, 1975, Voeller and Holt, 1973). Tree species that we and others (Southwick, 1975) have controlled with Tordon 101 Mixture are listed in Table 3. Maples should not be injected during heavy sap flow because of the washing action of sap.

Tordon 101 Mixture, due to freezing temperature susceptibility, is difficult to use in the winter. A new ready-to-use product now sold includes anti-freeze. We are now testing that product during the dormant (winter) season.

Tree injection gradually exposes "crop" trees to environmental forces, while felling unwanted trees with a chainsaw immediately opens up the stand. The exposed cambial layer of cut stumps still require painting or spraying. In either immediate or delayed felling, unwanted trees may be used later as firewood.

Table 3.—Tree species effectively controlled with injected Tordon 101 Mixture.¹

| SPECIES | SPECIES |
|----------------------|---------------------|
| American beech | Hickory spp. |
| American elm | Honeylocust |
| Big tooth aspen | Kentucky coffeetree |
| Big leaf maple | Mulberry spp. |
| Black birch | Osage orange |
| Black cherry | Pecan |
| Black locust | Persimmon |
| Black gum | Post oak |
| Black oak | Red bud |
| Black oak (Calif.) | Red elm |
| Blackjack oak | Red maple |
| Black walnut | Red oak |
| Black willow | Scarlet oak |
| Box elder | Service berry |
| Chestnut oak | Silver maple |
| Cottonwood | Southern red oak |
| Eastern hophorn bean | Sugar maple |
| Eastern red cedar | Sweetgum |
| Elm spp. | Sycamore |
| Flowering dogwood | Tan oak (Calif.) |
| Gray birch | Winged elm |
| Green ash | White birch |
| Hackberry | White oak |
| Hawthorn | |

1. Based on our study and those Southwick reviewed.

LITERATURE CITED

- Ferguson, Edwin R. and Edwin R. Lawson. 1975. Chemical injections for thinning pole-size hardwoods in the Ozarks. U.S. For. Ser. Res. Note SO-189. New Orleans, LA. 3 p.
- Geyer, Wayne A. and Larry E. Biles. 1975. Killing unwanted trees with Tordon 101 Mixture. Keeping Up With Research 21. Kan. Agr. Expt. Sta. Manhattan, K. 2p.
- Southwick, Lawrence, 1975. Tordon 101 R Forestry Herbicide. Indust. Veg. Mgmt. Vol. 7, No. 3. p. 14-17.
- Voeller, J. E. and H. A. Holt. 1973. Continued evaluation of the hypohatchet for woody species control. Proceedings 26th Annual Meeting, Southern Weed Science Society, p. 354-360.

Information in this report is for woodland owners, colleagues, industry cooperators, and other interested persons.

It is not a recommendation, but represents research at three locations.

Publications and public meetings by the Kansas Agricultural Experiment Station are available and open to the public regardless of race, color, national origin, sex or religion.