



2950 Niles Road, St. Joseph, MI 49085-9659, USA
269.429.0300 fax 269.429.3852 hq@asabe.org www.asabe.org

This is a peer-reviewed article.
2015 ASABE/IA Irrigation Symposium: Emerging
Technologies for Sustainable Irrigation
Proceedings of the 10-12 November 2015 Symposium,
Long Beach, California USA Published by
ASABE St. Joseph, Michigan, USA Publication date November 10, 2015
ASABE Publication No. 701P0415

An ASABE Meeting Presentation

(Accepted, Transactions of the ASABE)

Paper Number: 152145493

Cotton, Tomato, Corn, and Onion Production with Subsurface Drip Irrigation - A Review

Freddie R. Lamm

Kansas State University Northwest Research-Extension Center
Colby, Kansas, USA.
flamm@ksu.edu

**Written for presentation at the
Emerging Technologies for Sustainable Irrigation
A joint ASABE / IA Irrigation Symposium
Long Beach, California
November 10 – 12, 2015**

Abstract. *The usage of subsurface drip irrigation (SDI) has increased by 89% in the USA during the last ten years according to USDA-NASS estimates and over 93% of the SDI land area is located in just ten states. Combining public entity and private industry perceptions of SDI in these ten states, the major crops were tentatively identified as cotton, processing tomato, field corn, and onion. An extensive literature review of SDI usage for these four crops was performed concentrating on irrigation system comparisons, water and/or nutrient management, and SDI system design criteria. Although many crops potentially can be grown with SDI, the results presented here may be a relatively representative cross-section of the various opportunities and challenges of SDI for general crop production.*

Keywords. *microirrigation, SDI, trickle irrigation, irrigation methods, crop production.*

The authors are solely responsible for the content of this meeting presentation. The presentation does not necessarily reflect the official position of the American Society of Agricultural and Biological Engineers (ASABE), and its printing and distribution does not constitute an endorsement of views which may be expressed. Meeting presentations are not subject to the formal peer review process by ASABE editorial committees; therefore, they are not to be presented as refereed publications. Citation of this work should state that it is from an ASABE meeting paper. EXAMPLE: Author's Last Name, Initials. 2015. Title of presentation. In: Emerging Technologies for Sustainable Irrigation, ASABE / IA Irrigation Symp. St. Joseph, Mich.: ASABE. For information about securing permission to reprint or reproduce a meeting presentation, please contact ASABE at rutter@asabe.org or 269-932-7004 (2950 Niles Road, St. Joseph, MI 49085-9659 USA).

Introduction

Subsurface drip irrigation (SDI), the application of irrigation below the soil surface by microirrigation emitters (ASAE S526.3, 2007) is growing in usage in the USA and around the world. In the ten-year period (2003 to 2013), SDI in the USA increased by 89% from 164,017 to 310,361 ha (Fig. 1) according to USDA-NASS irrigation surveys, (USDA-NASS, 2004, 2010 and 2014). During the same time period SDI averaged approximately 25% of the surface drip irrigation (DI) land area. (Note: microsprinkler and bubbler irrigation are not included in these estimates). However, this comparison can perhaps be skewed by the fact that some of the SDI land area being reported is shallow, annually removed systems which are not intended for multiple-year use. The focus of this review is primarily on deeper systems intended for multiple-year usage.




Figure 1. Growth of land area in the USA using surface drip irrigation (DI) and subsurface drip irrigation (SDI), 2003 to 2013 according to USDA-NASS surveys (USDA-NASS, 2004, 2010, and 2014).

Subsurface drip irrigation has been the subject of three review articles during the past 20 years, Camp (1998) that provided an extensive characterization of the knowledge and studies that had been carried out to date, Rodriguez-Sinobas and Gil-Rodriguez (2012) that concentrated on design, uniformity, and soil water redistribution aspects, and Devasirvatham (2009) that focused on SDI for vegetable production. The status of the technology was also discussed in the 2000 and 2010 national irrigation symposiums sponsored by the American Society of Agricultural and Biological Engineers and The Irrigation Association (Camp et al., 2000 and Lamm et al., 2012). The goal of this review is to augment and supplement these earlier articles with a focus on the important SDI crops currently grown in the USA.

In 2013, the ten USA states with the largest SDI area (289,812 ha) comprise over 93% of the total SDI area but have a wide variation in the ratio of SDI/(SDI+DI) land area (Fig. 2). The variation can probably be explained by the crop production in these states with DI often being used on greater-value crops (typically fruits, nuts, and vegetables) and SDI being used on lesser-value commodity crops (e.g., corn, cotton, alfalfa, and other grain crops). There can be the persistent perception that SDI is harder to manage mainly because it has less visual cues that irrigation problems are occurring. As a result, many producers growing the greater-value crops choose DI as a less risky option and because cost of the irrigation system and its installation are not of paramount concern. When growing the lesser-value commodity crops with microirrigation, a deeper, multiple-year SDI system that can be amortized over several years is often the only economical option for a producer. This is particularly true in the Great Plains region where center pivot sprinkler irrigation has a good

economy of scale and the major sprinkler manufacturers are located (Nebraska). Although SDI land area in Nebraska and Kansas (near center of Great Plains) is relatively small (Fig. 2), the land area using SDI has grown 176 and 127% respectively in the last 5 years according to the USDA-NASS data. The principal irrigated crops in the Great Plains are the grain (primarily corn) forage (primarily alfalfa) and fiber crops (i.e., Cotton in Texas). However, there can be some exceptions to these overall SDI growth patterns. For example, California has the largest SDI land area in the USA and the primary crop using SDI is processing tomatoes. The growth of SDI in California has been 53% in the last 5 years (USDA-NASS, 2010 and 2014) and it is estimated that almost 40% of the irrigated cropland now uses some type of microirrigation (Taylor et al., 2014). No national or common state databases exists to link SDI land area with specific crop types, so only opinions from university, state and federal agencies, and industry personnel could be used to speculate on the crops utilizing the largest SDI land areas in the USA. Although, there is reasonable consensus that cotton, processing tomato, and field corn would constitute the largest three SDI usages, there is much less certainty about onion as being the fourth largest usage. Its inclusion in this review was most strongly supported by its reported prevalence in important SDI states in the Pacific Northwest (Oregon and Washington) and its presence in some of the other ten states. Although it is believed that total SDI vegetable land area excluding tomato and onion would exceed the SDI onion land area, totals of individual vegetable crops probably would not. It should also be noted that rotational crops grown on SDI land areas that are primarily for cotton, processing tomato, corn, and onion could add up to a considerable amount for some individual crop amounts, but these crops and land areas are unknown. There are vast geographical, topographical, and climatic differences across the USA that affect cropping and irrigation management. This review cannot exhaustively cover all of these aspects. The reader should consult individually reported studies for additional details.




Figure 2. Characteristics of SDI usage in the ten USA states having the largest land area devoted to SDI. The percentage of SDI refers to the ratio of SDI to total drip-irrigated land area (total SDI and surface drip irrigation (DI)). Data summarized from USDA-NASS (2014).

Cotton

Cotton (*Gossypium spp.*) production is one of the largest usages of SDI in the United States and also around the world. In suitable climatic regions compatible with cotton production, cotton with its lower water use can be an excellent crop in water short areas when coupled with efficient SDI. The largest usage of SDI for cotton in the United States is in Texas where 4263 SDI systems covering 107,356 ha primarily for cotton were reported in 2013 for the High Plains Water Conservation District alone (HPWCD, 2014). It can be noted that this regional value for Texas determined from SDI system physical locations is 55% greater than the overall Texas area shown in Figure 2. This illustrates the difficulty in finding good estimates for SDI land areas in the USA. The first reported use of SDI for cotton research that was found in this literature review was in 1963 near Lubbock, Texas (Zetzsche and Newman, 1996). In this one year trial, cotton lint yield was 0.80 and 0.77 Mg/ha for SDI and furrow irrigation with irrigation applications of 200 and 355 mm, respectively in this semi-arid region with summer pattern precipitation. Since that time, cotton has been the focus crop of many research studies and reports. Major SDI cotton research topics have been comparison of production under SDI and

alternative systems, response to SDI for a particular region, soil, or climate, water and nutrient management, and SDI system design criteria.

System comparisons for cotton

Sixteen studies were found in the literature that allowed equal or nearly equal comparisons of cotton lint yield under SDI and alternative irrigation systems (Table 1). Lint yield increases when using SDI as compared to alternative systems ranged from negative 1% to positive 65% with an average increase of 18% across all studies. Subsurface drip irrigated cotton lint yields were 2, 15, and 19% greater than DI, surface gravity, and sprinkler irrigation, respectively, in these studies. Greater incidence of spider mites with SDI as compared to sprinkler irrigation was reported in a study in California (Hollingsworth et al., 2014) which was attributed to drier and dustier leaves for the SDI system. In that same study, although yield differences were slightly greater for the sprinkler system and weed plant densities were similar, there was much greater weed biomass under the sprinkler. The most extensive SDI to sprinkler irrigation comparisons have been conducted in Texas (Bordovsky and Lyle, 1998 and Colaizzi, et al., 2010). In these studies SDI had 12 to 28% greater cotton lint yields when compared to three types of sprinkler application (Low elevation spray application, LESA; mid elevation spray application, MESA, and low energy precision application, LEPA), which are all considered to be efficient sprinkler application methods for this region. In these sixteen studies, the SDI lint yield advantage tended to be greatest when irrigation was limited and tended to decrease as irrigation levels increased and became more sufficient. This latter statement is generally consistent with results presented elsewhere for grain, fiber and forage crops (Camp, 1998, and Lamm et al., 2010) and possible reasons are that when water is sufficient, irrigation method matters less in overall yield and at greater irrigation levels there is increased possibility of nutrient leaching or soil aeration problems. Generally, the SDI increases in cotton lint yields as compared to alternative systems were greater on finer textured soils than coarse textured soils (Table 1). There can be notable exceptions to this conclusion, such as when lateral or upward soil water redistribution on heavier clay soils is negatively affected resulting in stand establishment or crop water stress later in the season. Soil water redistribution and aeration have been the topics of concern cotton production on clay soils in several SDI studies from Australia (Constable and Hodgson, 1990; Hulme and O'Brien 2000; Bhattarai et al., 2004 Bhattarai et al., 2005). On sandier soils, greater leaching may occur under SDI when irrigated with practices similar to other irrigation systems or when the sandy soil may not retain a large enough wetted volume for the seed establishment zone and crop root zone. However, 13% greater SDI lint yields as compared to furrow irrigation were reported on a nonuniform sandy loam soil with buried pockets of sand scattered throughout the field (Phene et al. 1992). The usage of SDI may have resulted in more consistent and uniform soil water within the crop root zone for this nonuniform soil even though the furrow irrigated treatment received 82% more irrigation. Crop water productivity with SDI was almost always greater in these seventeen system comparison studies, particularly in those studies where SDI was compared to surface gravity irrigation methods. These overall results might be partially attributed to greater partitioning of total water use to transpiration with SDI, less deep percolation, no irrigation runoff, and less evaporative losses (possibly both soil water evaporative losses and irrigation application losses). However, there is also growing evidence (Colaizzi et al., 2010) that SDI, by not wetting up the soil surface, is providing a more favorable thermal environment (warmer) and more heat units for irrigated cotton which is important, particularly as cotton production moves further northward in the Texas Panhandle, Oklahoma and southern Kansas.

Conjunctive nutrient and water management for cotton

Greater SDI cotton lint yield was obtained with an intermediate level of irrigation than with full irrigation on a sandy loam soil in Georgia (Sorensen, et al., 2004) which might be attributable to nitrogen leaching. They concluded that combined irrigation and nitrogen management with SDI might lead to decreased nitrogen requirements as compared to typical regional fertilization practices. A similar conclusion was stated for SDI cotton on a loamy sand soil for a four-year study in South Carolina that compared nitrogen application under every row and alternate row middle dripline spacings (Bauer et al., 1997, and Camp et al., 1997). The potential for nitrogen savings was similarly reported for field corn grown with SDI on a deep silt loam soil in Kansas (Lamm et al., 2001 and Lamm et al., 2004). Timing of nitrogen applications for cotton can affect yields and Constable et al. (1990) concluded that all N should be applied by mid flowering of the cotton and that perhaps as much as 30% of the total N should be applied before planting on a cracking clay soil in Australia. Excellent nitrogen and phosphorus uptake through SDI fertigation was reported for cotton resulting in greater N and P amounts in the leaf analyses in a study in Australia, but this did not result in greater yields (Aloni et al., 2000). The authors indicated that the yield results may have been confounded by differences in treatment water applications and dripline spacings. Extractable P concentrations in the immediate vicinity of a subsurface point source were 20 to 25% greater with continuous phosphorus fertigation than with periodic applications (Ben-Gal and Dudley, 2003) resulting in 20% greater biomass for sweet corn plants at 40 days post emergence.

SDI frequency for cotton

Frequent microirrigation events are often considered to be a necessary and desirable practice, but an SDI literature review (Camp, 1998) indicated that SDI frequency is typically only critical for shallow-rooted crops on shallow or sandy soils or in cases where salinity is a major concern. In a deficit-irrigated cotton study on a silty clay loam in Texas, frequency of SDI under deficit irrigation was not an important factor (Enciso-Medina et al., 2003). There were no differences in cotton

yield, quality, or gross returns related to SDI frequency, and the authors concluded that longer periods (as much as 16 days) between irrigation may allow irrigators to use less-expensive, manually operated SDI systems. Irrigation frequencies of 4 events/day, every two days, and weekly were compared under two irrigation levels (approximately 40 and 80% of ET) for cotton on a clay loam soil in Texas (Bordovsky, et al., 2012a). When averaged over the three years of the study, there were no statistical differences in yield for the drier irrigation treatment attributable to irrigation frequency and the every two day and weekly treatments were superior for the wetter irrigation treatment. A study in Israel on a heavy clay soil reported similar cotton seed yields for irrigation frequencies of twice or five times weekly (Plaut et al., 1985). These studies indicate high frequency is not required for cotton production. Cotton often benefits from some plant water stress in process of setting bolls and establishing overall plant size and it is possible irrigation frequency is less important because of this.

Dripline depth for cotton

Installation depth for SDI cotton has been a research topic in several studies (Kamara et al., 1991; Plaut et al., 1996; Khalilian et al., 2000; and Enciso et al., 2005). Cotton production was evaluated for dripline depths of 0.20, 0.31, and 0.41 m on a loamy sand with a clay hardpan at the 0.25 to 0.32 m depth in South Carolina (Khalilian et al., 2000). They found plant height was significantly greater for the 0.41 m depth and that the 0.20 m depth had greater weed infestations which required additional application of herbicides. Lint yields were greater with increased depth in both years of the study averaging 1.29, 1.38, and 1.47 Mg/ha for dripline depths of 0.20, 0.31, and 0.41 m, respectively. In a study in semi-arid Lubbock Texas (USA) on a fine sandy loam soil, cotton root development and distribution were not affected by dripline depths of 0, 0.15, 0.30 and 0.45 m (Kamara et al., 1991). Their research suggests that in regions that typically receive precipitation during the active growing season, that dripline depth will not be the overriding factor in root development and distribution. In a laboratory column study on a loessial brown loam using SDI, Plaut et al. (1996) found that cotton roots can develop under partial wetting of the upper soil profile to soil water potentials of 0.1 MPa. They suggested that a reasonable dripline depth for cotton would be 0.4 to 0.5 m. Lint yields were 5% greater for a 0.3 m dripline depth as compared to a 0.2 m depth on a clay loam site in western Texas (Enciso et al., 2005). In the sixteen studies listed in Table 1, the average depth was 0.32 m. In summarizing these overall results, deeper dripline depth (0.30 m or greater) tends to be advantageous for cotton provided that germination/crop establishment are not a concern.

Dripline spacing for cotton

Dripline spacing for cotton has been the focus of SDI research studies in Georgia (Sorensen et al., 2011), South Carolina (Bauer et al., 1997, Camp et al., 1997, and Khalilian et al., 2000) and Texas (Enciso-Medina et al., 2005). In the humid region (South Carolina), a dripline spacing of 2 m for cotton using an alternate row/bed middle pattern (i.e., one dripline centered between two rows) was usually adequate and more economical, saving about 30% in SDI system costs compared to a 1 m dripline spacing (Bauer et al., 1997, and Camp et al., 1997). Averaging eight years of lint yield data from SDI on a sandy loam in Georgia, Sorensen et al. (2011) concluded an alternate row furrow lateral spacing was as effective as driplines placed under every crop row. Similarly, no cotton lint yield differences were reported for every row vs alternate row-middle dripline placements on a loamy sand in another study from South Carolina (Khalilian et al., 2000). The wider alternate row-middle spacings might present greater germination and crop establishment problems during drought periods on coarser textured soils as suggested by Camp et al. (1997). Similarly, a decreased SDI wetting pattern effects can occur on hardsetting soils with poor soil structure. It was concluded that a wider 2 m dripline spacing, commonly used for cotton in Australia was not appropriate on hardsetting red soils due to the limited flow perpendicular to the dripline (Hulme and O'Brien, 2000). In a three year study in semi-arid Texas, overall cotton lint yields were 21% greater for a 1 m dripline spacing placed under every crop row than for an alternate row-middle dripline spacing of 2 m (Enciso et al., 2005). In this study, yields were severely reduced for the wider spacing in one year due to seed germination failure. However, in one year, the wider spacing outperformed the narrower dripline spacing leading the authors to conclude that further research was needed with regard to dripline placement and its interaction with wheel traffic and tillage/bed management. The climate, soil, and crop rooting characteristics can affect the required dripline spacing and placement. When germination and crop establishment is less of a concern, there is general agreement that the alternate row/bed dripline spacing (about 1.5 m) is adequate for most of the deeper-rooted agronomic crops such as cotton on medium- to heavy-textured soils (Camp, 1998).

Summary of SDI for cotton

Cotton lint yield increases averaging 19% as compared to sprinkler irrigation as listed here are illustrative of why SDI is increasing being adopted in Texas. The yield increases tended to be greatest for SDI when irrigation was limited. Conjunctive management of nutrients and irrigation should help further SDI adoption for cotton. Frequency of SDI has not been a major issue and this has allowed some producers to use less costly, manually-operated systems. Dripline depths of approximately 0.30 m allows for normal cultural practices and allow for long term, multiple-year SDI systems. Dripline spacings are most often an alternate row middle arrangement which helps to keep SDI system costs down. Cotton is a major commodity crop in the USA, but does not have the net returns that are possible with the greater value fruits and vegetables. Therefore, reliable, multiple-year, and cost-effective SDI systems are important for cotton.

Table 1. Cotton lint yield, Mg/ha as affected by irrigation method and average lint yield increase attributable to SDI.

Research report	Additional factors or comments ¹	Soil type	SDI dripline depth (m)	Subsurface drip (SDI)	Surface drip (DI)	Surface gravity	Other sprinkler	MESA sprinkler	LESA sprinkler	LEPA sprinkler	Average increased yield using SDI over all alternative methods, %
Ayars et al., 1999	All years	Silty clay	0.4 and 0.6	1.73	-	1.47	-	-	-	-	18
Bordovsky and Lyle, 1998	2.5 mm/d capacity	Loam	0.3	1.28	-	-	-	-	-	1.05	22
	5.1 mm/d capacity			1.37	-	-	-	-	-	1.26	9
	7.6 mm/d capacity			1.41	-	-	-	-	-	1.31	8
	LEPA frequency, 1 d			1.36	-	-	-	-	-	1.20	13
	LEPA frequency, 2 d			1.36	-	-	-	-	-	1.21	12
	LEPA frequency, 3 d			1.36	-	-	-	-	-	1.20	13
Bordovsky and Porter, 2003	Limited preseason irrigation	Loam	0.3	1.14	-	-	-	-	0.83	0.95	28
	Full preseason irrigation			1.18	-	-	-	-	0.95	1.05	18
	2.5 mm/d capacity			1.01	-	-	-	-	0.71	0.85	29
	5.1 mm/d capacity			1.31	-	-	-	-	1.07	1.14	18
Bhattarai et al., 2005	SDI data for 75% ET	Clay	0.4	1.93	-	2.15	-	-	-	-	-10
Colaizzi et al., 2010	25% of full irrigation	Clay loam	0.3	0.64	-	-	-	0.46	0.49	0.55	28
	50% of full irrigation			0.80	-	-	-	0.56	0.56	0.74	29
	75% of full irrigation			1.02	-	-	-	0.78	0.75	0.87	28
	100% of full irrigation			1.07	-	-	-	0.87	0.89	0.99	16
Constable and Hodgson, 1990	All varieties	Clay	0.2	1.81	1.77	1.76	-	-	-	-	3
Grabow et al., 2002	-	Clay loam	0.3 and 0.45	1.42	-	-	1.21	-	-	-	17
Hanson, et al., 1970	Heavy irrigation	Not listed	0.3	1.62	-	1.41	-	-	-	-	15
	Light irrigation			1.53	-	1.44	-	-	-	-	6
Hollingsworth et al., 2014	-	Clay loam	0.3	1.46	-	-	1.48	-	-	-	-1
Nuti et al., 2006	-	Sandy loam	0.25	1.47	-	-	1.40	-	-	-	5
Phene et al., 1992	Good soil	Sandy loam	0.375	1.87	-	1.89	-	-	-	-	-1
	Poor soil			1.78	-	1.56	-	-	-	-	14
Smith et al., 1991	-	Fine textured	0.45	1.71	-	1.20	-	-	-	1.14	46
Sorensen et al., 2004	2000 data for full irrigation	Sandy loam	0.30 to 0.35	1.26	-	-	1.26	-	-	-	0
Thompson and Enciso, 2002	Commercial farm data	NA	NA	1.05	-	0.71	-	-	-	-	48
Tollefson, 1985	SDI and furrow	NA	0.20 to 0.25	2.12	-	1.60	-	-	-	-	33
	SDI and sprinkler			2.23	-	-	1.35	-	-	-	65
Whitaker et al., 2008	Data for two locations	Loamy sand	0.25	1.41	-	1.32	-	-	-	-	6
Average SDI increase across all studies and treatments											18
Average SDI increase over surface gravity irrigation across all studies											15
Average SDI increase over all sprinkler irrigation treatments across all studies											19
Average SDI increase over DI across all studies											2

Tomato

Tomato (*Solanum lycopersicum* L.) production is one of the largest usages of SDI in the United States, primarily in California for processing tomato. Estimates from University of California-Davis (Montazar, 2015) indicate that SDI has become the commercial standard for processing tomato with over 100,000 hectares or approximately 80% of the

processing tomato land area. This would suggest that 60 to 65% of the SDI land area in California is used for processing tomato. The advantages of using SDI for processing tomatoes was thoroughly discussed by Phene et al., 1987. Much of this early work by Phene et al. and follow-up research was summarized by Ayars et al., 1999. Tomato has been the focus crop of many other research studies and reports. Major SDI tomato research topics have been comparison of production under SDI and alternative systems, response to SDI for a particular region, soil, or climate, water and nutrient management, and SDI system design criteria.

System comparisons for tomato

A total of sixteen studies were found in the literature that allowed equal or nearly equal comparisons of tomato yield under SDI and alternative irrigation systems (Table 2) with SDI yield increases ranging from negative 32% to positive 205% with an average positive increase of 12% across all studies. Subsurface drip irrigated tomato yields were 7, 17, and 23% greater than DI, surface gravity, and sprinkler irrigation, respectively, in these studies. Large yield reductions (31 to 32%) with SDI were reported in three studies for which the authors suggested possible rationale. A yield reduction (32%) with SDI was reported on a fine sand in Florida and it was suggested yield reductions were the result of water availability rather than nutrient management (Clark et al., 1993). The shallot zone of tomato was insufficient to efficiently use irrigation water applied by driplines at a 0.3 m depth. In a study in California on a clay loam soil examining the effects of SDI on weed control, a yield reduction of 32% was attributed to early cutoff of irrigation for the SDI treatment (Shrestha et al., 2007). Although SDI yields were severely affected by irrigation management, the authors did report that weed emergence was nearly eliminated in the furrows and the amount of weeds on the crop bed was reduced from 46 to 96%. Similar reports of reduced weed growth with SDI as compared to sprinkler irrigation was reported for processing tomato by Grattan et al. (1990). Greater processing-tomato yield with DI than SDI was attributed to serious root intrusion that occurred mid-season in SDI for a study in Canada (Tan et al., 2003). It was suggested that water stress that likely occurred before the SDI system was replaced with surface drip for the remainder of the season caused the 31% yield reduction. In contrast, more than one-half of the studies had tomato yield increases of 10% or greater for the most productive treatments (Table 2).

Conjunctive nutrient and water management for tomato

The studies conducted by the USDA Water Management Research Laboratory in California (Ayars et al., 1999) are worth noting in that they illustrate how combined irrigation and nutrient management can be so beneficial to tomato production with SDI. Tomato yields were 10% greater on a clay loam soil when phosphorus (P) and potassium (K) were injected with high frequency SDI compared to low frequency DI, without a concurrent increase in crop water use, thus resulting in 13% greater crop water productivity for the high frequency SDI (Phene et al., 1990). Rooting and nutrient uptake patterns were deeper with SDI and P fertigation, and similarly, petiole P deficiencies were observed when P was not injected (Ayars et al., 1999). Tomato yields were 35% greater with daily SDI than with events every third day on a calcareous soil in Egypt (El-Gindy and El-Araby, 1966). The more frequent irrigation events resulted in wider wetting patterns and may have reduced salinity in the vicinity of the roots. However, Hanson et al. (2003) reported no yield benefit to SDI (0.2 m depth) for multiple irrigation events per day for processing tomato compared with frequencies as low as weekly on a silt loam soil in California. The phosphorous fertilizer in this study was applied preplant, and the nitrogen was applied weekly. These studies show that irrigation frequency can interact with soil characteristics and nutrient management practices in different ways.

Dripline depth for tomato

Various dripline depths have been compared in a few studies for SDI tomato production (Schwankl et al., 1990; Machado et al., 2003 and 2005). Seed germination of direct-seeded processing tomato at a seeding depth of 0.038 m averaged 219% better at dripline depths of 0.15 and 0.23 m than at 0.30 m (Schwankl et al., 1990) when averaged over three seedling irrigation rates on a Yolo clay loam soil in California. The authors concluded that acceptable tomato stands could be achieved with SDI as the sole source of irrigation for germination. In some areas, sprinkler irrigation is used for tomato crop establishment, so using SDI for this purpose would be less of a concern. No significant differences were reported for dripline depths of 0.2 and 0.4 m for tomato production on sand for two studies in Portugal (Machado et al., 2003 and 2005) although they did report conflicting results for SDI and DI comparisons (Table 1). In the sixteen studies (Table 2), the average depth was 0.27 m with a dripline depth range of 0.15 to 0.25 m being most common. There has been increased use of transplants for processing tomato in California and the transplanting process would probably not rely solely on SDI for crop establishment.

Salinity management for tomato

Soil and water source salinity can be a constraint in tomato production and SDI can potentially help manage salinity. Little correlation was found for processing-tomato yield and soil salinity level for SDI in a study in California (Hanson and May, 2003a). They concluded that soil salinity under SDI may affect tomato yield less than with other irrigation methods. This could be because SDI is providing a more consistent soil water environment in the soil than alternative systems. Tomato yields in this study were 12.9 to 22.6 Mg/ha greater with SDI than with sprinkler irrigation for similar irrigation amounts. Sodium and chloride accumulations were reduced in tomato plant tissues with SDI compared to DI on

a silty clay soil in Tunisia, resulting in greater leaf area and chlorophyll for the SDI tomato plants (Kahlaoui et al., 2011). Less salt accumulation was present in the active root zone and denser root systems occurred with SDI than with DI for tomato on a calcareous soil in Egypt (El-Gindy and El-Araby, 1996). This was attributed to better soil water redistribution within the root zone. Greater horizontal movement of the water occurred with larger 4 L/h emitters compared with greater downward movement for 2 L/h emitters. However, there was still an expressed concern that SDI may cause an excessive salt accumulation in the seed and transplant zone for the next crop. The management of crop location with respect to dripline location can be important for even moderately saline waters with SDI systems that are used for multiple years unless periodic leaching is provided. Root activity was limited to the wetted soil volume for drip-irrigated tomato on a sandy soil, but the rooting patterns were different for fresh and saline water (Ben-Asher and Silberbush, 1992). When freshwater was used, a relatively high root density occurred around the periphery of the wetted volume, but with saline water limited root activity existed at the periphery. In summarizing the studies listed here, it appears that with precise management SDI can be appreciably better than other irrigation systems for use with saline waters for tomato production.

Summary of SDI for tomato

Yield increases averaging 12% as compared to alternative irrigation systems as listed here are illustrative of why SDI has become the commercial standard for processing tomato with large SDI land areas in California. Combined nutrient and water management which is easily attainable with SDI results in sizable yield increases. A dripline depth range of 0.15 to 0.35 m is common. Salinity management for tomato production in arid regions can be enhanced with SDI, but care must be employed to avoid salinity hotspots within the crop beds.

Table 2. Tomato yield, Mg/ha as affected by irrigation method and average yield increase attributable to SDI.

Research report	Substudy or comments ¹	Additional factors	Soil type	SDI dripline depth (m)	Subsurface drip (SDI)	Surface drip (DI)	Surface gravity	Sprinkler	Average increased yield using SDI over alternative methods, %
Ayars et al., 1999	CSUF study, 1981	-	Sandy loam	0.46	132	-	90	-	47
	Westside Field Station study, 1984-87. Hi Frequency SDI & DI	N-Only, 1984	Clay loam	0.45	121	126	-	-	-4
		N + P, 1985			168	152	-	-	11
		N + P + K, 1987			220	201	-	-	9
	Westside Field Station study, 1987. Hi Frequency SDI & DI Continuous P injection	No P	Clay loam	0.45	184	177	-	-	4
		P, 67 kg/ha			220	201	-	-	9
		P. 134 kg/ha			215	192	-	-	12
	Westside Field Station study, 1987. Hi Frequency SDI & DI	N-Only, 1990	Clay loam	0.45	143	143	-	-	0
		N + P, 1990			182	155	-	-	17
		N + P + K, 1990			179	154	-	-	16
Batchelor et al., 1994	Small garden study	Avg. of 6 irrigation trts.	Sandy clay loam	0.15	96	-	96	-	0
Bogle et al., 1989	Fresh market tomatoes	Spring 1983	Sandy clay loam	0.15 to 0.20	39	-	32	-	22
		Fall 1983			34	-	28	-	21
		Spring 1984			52	-	43	-	21
Clark et al., 1993	Fresh market tomatoes	Spring 1993	Fine sand	0.30	54	79	-	-	-32
del Amor and del Amor, 2007	Two irrigation levels	50% irrigation	Clay loam	0.40	58	19	-	-	205
		100% irrigation			76	70	-	-	9
El-Gindy and El-Araby, 1996	Varying irrigation frequency and emitter discharge on high salinity soil	Daily, Hi Flow	Loamy mixed	0.25 to 0.30	45	35	-	-	29
		Daily, Lo Flow			33	33	-	-	0
		3 day, Hi Flow			30	34	-	-	-12
		3 day, Lo Flow			28	31	-	-	-10
Grattan et al., 1990	Weed study for processing tomato, reds shown here.	No Herbicide	NA	0.25	119	-	78	78	53
		Herbicide			117	-	101	105	14
Hanson and May, 2000	Processing tomato	Year 1999 data for 2 sites	Silty clay	0.20 to 0.23	104	-	-	82	27
			Clay loam		91	-	-	79	15
Hanson and May, 2003a	Processing tomato	Year 2000 data	NA	0.20	86	-	-	64	34
Hanson and May, 2007	Processing tomato	Years 2004-2006	NA	0.23	114	96			18
Machado et al., 2003	Average of 1997- 1998	0.2 m depth SDI	Sand	See trt. depth	130	115	-	-	13
		0.4 m depth SDI			128	115	-	-	11
Machado et al., 2005	Averaged across 3 irrigation levels	0.2 m depth SDI	Sand	See trt. depth	118	119	-	-	-1
		0.4 m depth SDI			107	119	-	-	-10
Marouelli and Silva, 2002		0.2 m depth SDI	Clay	See trt. depth	116	124		108	0
		0.4 m depth SDI			94	124		108	-28
Najafi and Tabatabaei, 2007		0.15 m depth SDI	Silty loam	See trt. depth	52	53	44		8
		0.30 m depth SDI			37	53	44		-23
Shrestha et al., 2007	Conventional vs. conservation tillage	2004 and 2005 data	Clay loam	0.23	64	-	94	-	-32
Tan et al., 2003	Study on two soil types	Broadcast fertilization	Loamy sand	NA	139	130	-	-	7
		Fertigated			139	133	-	-	5
	<i>SDI failed due to root intrusion and was replaced with DI inseason for clay loam</i>	Broadcast fertilization	Clay loam	NA	97	114	-	-	-15
		Fertigated			84	121	-	-	-31

Average SDI increase across all studies and treatments	12
Average SDI increase over surface gravity irrigation across all studies	17
Average SDI increase over all sprinkler irrigation treatments across all studies	23
Average SDI increase over DI across all studies	7

Corn

Field corn (*Zea mays* L.) is the largest irrigated crop in the USA with typically 4.85 to 5.40 million hectares harvested for grain in recent years (USDA-NASS, 2014), so even though the land area using SDI is much smaller than for cotton and tomato production it is still an important usage and has been the topic of many research studies. Although, the SDI land area for corn could not be determined, university and government agency anecdotal responses to inquiries by this author in early 2015 reported some SDI corn production (grain and/or forage) in six (Florida, Georgia, Kansas, Nebraska, Oregon, and Texas) of the top ten states having the largest SDI land area in the USA (USDA-NASS, 2014). The earliest reported usage of SDI for corn in the USA was in 1915-1916 on a deep silt loam soil at the West College Farm of Colorado State University (House, 1918). The description of the irrigation system is somewhat similar to a subirrigation system, but this system was not used for controlling the height of the water table. In this study, porous clay tiles were installed at a depth of 0.6 m on various lateral spacings of 4.9 to 7.6 m. Precipitation in 1915 at the location was greater than normal, so corn yield measurements were not recorded. Dry conditions in 1916 resulted in large plant height differences ranging from approximately 1 to 2 m across the width of the widely-spaced subsurface laterals. They concluded that the technology would not be cost effective for ordinary farm crops such as corn, that narrower lateral spacings would be needed for grain crops on this soil type, and that a smaller water supply that would be infeasible for a surface gravity system could be sufficient for a subsurface system. In a three-year study (1965-1967) near Georgetown, Delaware, Mitchell et al. (1969) reported some early SDI corn research evaluating flexible plastic tubing with various orifices and dripline spacings (1, 1.5, and 2 m). They reported yield increases of 12 to 2515% with SDI compared to rainfed production on a loamy sand. This report also provides some of the earliest details about installation implements and procedures for SDI, some of which are similar to today's procedures. In another related early publication, Mitchell and Tilmon (1982) suggested SDI as a good, economical, irrigation system alternative for the small farmer in the USA. This is because the components of SDI systems can be easily and economically designed to accommodate the field size (Bosch et al., 1992; O'Brien et al., 1998). Several early SDI corn studies (1989 to 1999) from Kansas were discussed by Lamm and Trooien (2003). Major SDI corn research topics have been comparison of production under SDI and alternative systems, response to SDI for a particular region, soil, or climate, water and nutrient management, and SDI system design criteria.

System comparisons for corn

A total of twelve studies were found in the literature that allowed equal or nearly equal comparisons of corn grain yield under SDI and alternative irrigation systems (Table 3) with SDI yield increases ranging from negative 51% to positive 30% with an average positive increase of 4% across all studies. Subsurface drip irrigated corn grain yields were 7, -16, and 0% greater than DI, surface gravity, and sprinkler irrigation, respectively, in these studies. In a study in Egypt, large grain yield declines (23 to 51%) associated with SDI as compared to furrow irrigation and DI were attributed to the reduced ability of SDI to manage salinity on the loamy soil (Abou Kheira, 2009). In contrast in another study from Egypt (Abuarab et al., 2013), a large yield increase occurred with SDI as compared to DI (23 and 38% with and without SDI air injection, respectively) in a two-year study on a sandy clay loam. Corn grain yield varied widely by annual weather conditions for SDI and simulated LEPA sprinkler irrigation in a seven-year study on a deep silt loam soil in Kansas (Lamm, 2004). Although on the average, yields were similar between the two systems, somewhat surprisingly, yields were approximately 5% greater with SDI in normal to wetter years and approximately 4% greater for LEPA sprinkler in severe drought years (Table 1). Greater LEPA yields were associated with greater kernels/ear in the drought years (LEPA, 534 vs. SDI 493 kernels/ear), while SDI had greater kernel mass in all years (SDI, 347 vs LEPA, 332 mg/kernel). Although, it appears there is little evidence to suggest SDI greatly increases yields over other well-managed irrigation systems, Lamm (2014) suggests that under deficit irrigation, SDI may stabilize corn yields at a greater level than center pivot sprinkler irrigation. Water productivity was generally greater for SDI in most of the studies that might be attributable to less soil water evaporation, less deep percolation, and reduced irrigation and precipitation runoff. Subsurface drip irrigation can economically compete with less expensive center pivot sprinkler systems by increasing irrigated land area (i.e., no unirrigated corners as is often the case with center pivot sprinklers), particularly when corn selling prices and grain yields are high (O'Brien et al., 1998 and Lamm et al., 2015).

SDI frequency for corn

Irrigation frequencies ranging from daily to weekly events have had very little effect on field corn production, provided soil water was managed within acceptable stress ranges (Caldwell et al., 1994; Camp, et al., 1989; Howell et al., 1997). Irrigation water use efficiency was increased 16% by decreasing the frequency from one to seven days and was attributed to improved storage of precipitation on a deep silt loam soil in the semiarid region of Kansas (Caldwell et al., 1994). In a

later Kansas corn study using a deficit irrigation capacity of 3.8 mm/d, Lamm and Aiken (2005) also found no major effect of SDI frequencies ranging from one to seven days. Irrigation frequency had an effect in only one of the three years of the study, and in that extreme drought year, the less-frequent, seven-day treatment had higher grain yield. The number of kernels/ear was greater for the less-frequent irrigation treatments in this drought year. It was hypothesized that the larger irrigation amount (i.e., 26.7 mm/7 d) for the less-frequent treatment established a larger wetted root zone, allowing for better early season ear and kernel development. All of these studies were conducted on medium to heavy textured soils, so greater frequency could still be important on lighter textured soils.

Dripline depth for corn

Long term, multiple-year SDI systems are important when growing lesser value commodity crops such as corn, so deeper dripline depths can be advantageous in avoiding mechanical damage due to tillage and also possibly pest damage (Lamm and Camp, 2007). Dripline depth for SDI corn has been the experimental factor in four research studies found in the literature. Corn grain yields were unaffected by dripline depths ranging from 0.2 to 0.4 m, and there was only a slight yield reduction at the 0.5 and 0.6 m dripline depths on a deep silt loam soil in Kansas (Lamm and Trooien, 2005). They concluded that factors external to the study (e. g., producer preferences, tillage schemes, rodent management, need for surface wetting for crop establishment, installation draft requirements and cost) might have a larger influence on selection of dripline depth. Dripline depths of 0.15, 0.23 and 0.3 m were evaluated under two alternative crop/dripline placements (standard design with driplines spaced at 1.5 m beneath the crop furrow, and wide twin-row crop bed design with 1.5 m dripline spacing) for corn in a Texas study at Bushland, Texas on a clay loam soil (Colaizzi et al., 2006 and Bordovsky et al., 2012b). Although favorable weather conditions negatively affected the determination of dripline depth effects on corn germination and establishment during the three year period (2006 to 2008), the authors reported on average corn grain yields were greatest for the standard design when dripline depth was 0.23 m and when the dripline depth was 0.30 m for the wide twin-row crop bed. There were no statistically significant differences in dripline depths of 0.25, 0.35 and 0.45 m for forage corn production on a loamy soil in Mexico (Montemayor Trejo et al., 2006) although water productivity for all three dripline depths were 45% greater than for a furrow-irrigated treatment. However, in an SDI corn study on a sandy loam soil in New Mexico, greater plant stands, greater yields, and greater irrigation water use efficiency was reported for 0.15 and 0.2 m dripline depths than for 0.25 and 0.30 m depths (Pablo, et al., 2007). Plant stands were reported averaging 98% for the 0.15 m depth to a dismal 50% for the 0.3 m depth in the two-year study, with the authors attributing much of the yield differences to poor crop stands for the deeper depths. Sweet corn yields were 31% greater for dripline depths of 0.23 m as compared to 0.33 m on well-drained sandy soil in Florida (Dukes and Scholberg, 2005). They also reported up to 24% less deep percolation with SDI as compared to sprinkler irrigation. Annual soil water evaporation loss reductions of 23 and 64% were predicted for 0.15 and 0.30 m dripline depths, respectively, compared with DI in a field and modeling study for corn on a Pullman clay loam soil in Texas (Evelt et al., 1995), while having no appreciable effect on corn transpiration. Although deeper dripline placement will minimize soil water evaporation losses, this aspect must be balanced with the potential for increased percolation losses, while considering the crop root-zone depth and rooting intensity (Gilley and Allred, 1974; Thomas et al., 1974; Philip, 1991). Deeper dripline depths can also be beneficial in weed control for SDI field and sweet corn as reported by Bar-Yosef et al., (1989), Oron et al., (1991b) and Lamm and Trooien, (2005). In a study from Israel with a dripline depth of 0.3 m on a loessial soil, weed growth and the resultant nitrogen (N) uptake by the weeds was reduced by almost 28% compared with DI for sweet corn production (Bar-Yosef et al., 1989). In the twelve studies listed in Table 3, the average dripline depth was 0.29 m with depths ranging from 0.15 to 0.45 m. This range of values compare well with the range reported for corn in the SDI review by Camp (1998). In summarizing these dripline depth results for corn, it can be said that both field and sweet corn have extensive root systems that can tap into water applied with deeper driplines (approximately 0.3 to 0.4 m), that crop germination may be improved by shallower driplines (approximately 0.2 to 0.25 m) in areas where crop establishment is a concern, and that retaining the applied water in the crop root zone is important (i.e., minimizing soil water evaporation and deep percolation) in obtaining the benefits of SDI.

Dripline spacing for corn

Dripline spacing greatly affects SDI system costs, so it should not be surprising that it has been the focus of several studies for a lesser-value commodity crop such as corn. As a general rule, SDI dripline spacing is an integer multiple of the crop row spacing and the crop row or bed spacing is usually set by cultural practices for a given crop in a given region. Dripline spacing for corn is typically an alternate row/bed middle pattern with one dripline per bed or between two rows, but in a few cases might be one dripline per row/bed. It appears more dripline spacings studies for corn have been conducted in the humid and semi-humid regions than in the semi-arid and arid regions. This is probably because the annual profitability of irrigation varies more in these wetter regions and researchers are looking for possible practical ways to reduce irrigation investment costs. Grain yield for a 1 m dripline spacing was 10% and 28% greater than for 1.5 and 2.0 m spacings for corn grown with a 0.9 m row spacing for a loamy sand in Delaware (Mitchell et al., 1969). However, the 2.m dripline spacing still increased the yield over rainfed production by 18%. Average corn yields were 100, 93, and 94% of the maximum yield for dripline spacing of 0.91, 1.83, and 2.74 m in the humid region of Virginia under adequate irrigation on a loamy sand (Powell and Wright, 1993). In an economic comparison of these results, it was concluded that the 1.8 m spacing was best for corn and peanut rotations (Bosch et al., 1998.) for Virginia and North Carolina. An every-

crop row dripline spacing as compared to an alternate-row middle placement resulted in greater corn grain yields in 50% of the years in an eight year study on a sandy loam soil in Georgia (Sorensen et al., 2013). However, because the narrower dripline spacing only resulted in 4% greater yields (9.2 vs 8.8 Mg/ha), the authors concluded that the narrower dripline spacing would probably not be economically justified. Greater row-to-row variability was reported for narrow-row spaced corn (0.38 m) for wider dripline spacings (2 vs. 1 m) on a loamy sand in South Carolina, leading the authors to suggest that increasing planting density nearer the driplines might improve productivity (Stone et al., 2008). The 1 m dripline spacing increased grain yields by 12% compared to the 2.m spacing (Table 3). Similar corn grain yields were reported for 0.76 and 1.52 m dripline spacings on a clay loam soil in North Carolina (Grabow, et al., 2002). Corn grain yields were 13.6, 12.9, 12.3, and 11.7 Mg/ha for dripline spacings of 0.76, 1.52, 2.28 and 3.04 m, respectively, for 0.76 m spaced corn rows that were planted perpendicular to the driplines on a silt loam soil in Kansas (Spurgeon et al., 1991). Although yields were greater for the 0.76 m dripline spacing in this three-year study, the authors concluded the additional dripline costs would not be justified. Dripline spacing that is too wide can lead to excessive deep percolation below the crop root zone (Darusman et al., 1997), however, and large yield reduction or crop failure in rows that are furthest from the dripline (Lamm et al., 1997). Although SDI soil water redistribution is often correlated to some degree to soil texture (i.e., percentages of sand, silt, and clay), wetting can be very different for layered soils of similar texture (Cote et al., 2003; Thorburn et al., 2003). Soil water redistribution models are seldom used for determination of optimal dripline spacing because of the domination of grower preferences in crop row/bed configurations and their interrelationship with dripline spacing. In summarizing the dripline spacing results for corn, it can be said wider dripline spacing may be suitable in soils with layering, allowing increased horizontal soil water redistribution above the soil layer, and in regions that are less dependent on irrigation for crop production, but overall an alternate row/bed dripline spacing (about 1.5 m) is usually adequate for corn on medium- to heavy-textured soils. These results match the conclusions of the review by Camp (1998) and the historical discussion by Camp et al. (2000).

Conjunctive nutrient and water management for corn

Fertigation with SDI can be environmentally beneficial as well as potentially improving corn yields through improved plant nutrition and root growth. Some forms of N are readily leachable, so SDI can be a good tool for timely applications with precise placement in the crop root zone. Combined management of irrigation and anticipated rainfall has long been a necessary tool to manage nitrogen fertilization on sandy soils. Subsurface drip irrigation of water containing urea-ammonium nitrate (UAN, 32-0-0) can supply both the mobile, but readily absorbed nitrate-N and the less mobile ammonium-N, which can be absorbed directly by the plant or microbially transformed to nitrate-N. In a Kansas study on a deep silt loam soil, corn grain yield, plant nitrogen uptake, and water productivity were statistically unaffected by the N application method (preplant surface application or in-season SDI fertigation), but all three factors were influenced by the combined total N amount (Lamm et al., 2001). Nearly all of the residual nitrate-N after corn harvest was within the upper 0.3 m of the soil profile when all the N was preplant-applied N, regardless of irrigation regime (75, 100, and 125% of ETc). In contrast under SDI fertigation, nitrate-N concentrations increased with increasing rates of injected N and migrated deeper into the soil profile with increased irrigation. The results suggest that total N for corn production potentially could be reduced when using SDI fertigation as compared to surface-applied preplant N banded in the furrow. In a follow-up study at the same location (Lamm et al., 2004), a best management practice (BMP) for SDI nitrogen fertigation of corn was developed using the criteria of residual ammonium- and nitrate-N levels in the soil profile, grain yield, plant nitrogen uptake, and water productivity. The overall BMP included scheduling irrigation to replace 75% of the calculated soil water deficit attributable to evapotranspiration, and weekly SDI N fertigations, with the total in-season N fertigation amount not to exceed 180 kg/ha and the total applied N not to exceed 210 kg/ha. A single early season N application was compared to multiple in-season SDI fertigation events on a silt loam soil on Nebraska (Tarkalson and Payero, 2008). Averaged over three years and three total seasonal N rates (128, 186, or 278 kg N/ha), corn grain yield, plant biomass, and plant nitrogen uptake were 4, 1, and 10% greater with the multiple in-season SDI fertigation events. Greater N use efficiency when using SDI fertigation compared to surface N applications was also reported for corn production on a silt loam soil in Delaware (Mitchell, 1981) with greater P and K availability mentioned as well. Several SDI studies (Bar-Yosef et al., 1989; Martinez-Hernandez et al., 1991; Phene et al., 1991, and Ben-Gal and Dudley, 2003) have examined the effect of phosphorus fertigation on sweet corn production and their results may suggest new research for field corn is needed as both irrigation and nutrient management become more intensively managed. Fertigation of P through SDI increased marketable ear yield by 12% compared to DI and moved the center of the root density to a deeper depth (i.e., 0.3 m for SDI and 0.1 m for DI) on a loessial soil under net/screen house production in Israel (Martinez-Hernandez et al., 1991). Similar results were reported in a field study in California on a clay loam soil for sweet corn P fertigation by Phene et al. (1991) who found that DI had the greatest root density in the 0 to 0.3 m depth, whereas SDI had greater root density below 0.3 m. Sweet corn yield increases of 4 to 10% were reported on a loessial soil in Israel for SDI when compared with DI (Bar-Yosef et al., 1989). Although, total biomass P uptake was unaffected by irrigation system type, the increased ear yield response with SDI was attributed to greater dry matter allocation to the ear. Although P is considered a relatively, immobile nutrient, frequent SDI fertigation can potentially move P under mass flow to a larger portion of the active corn root zone (Martinez-Hernandez et al., 1991 and Ben-Gal and Dudley, 2003). In summarizing the SDI fertigation studies for corn, it can be said that SDI can be an excellent irrigation system for improved nutrient management. The results emphasize that high-yielding corn production also can be environmentally sound and efficient in

both nutrient and water use. However a caveat is worth mentioning. When water and nutrients are highly managed for greatest effectiveness, there can be less margin of error. It is important that producers are diligent in observing the corn growth and development. Under drought conditions, preplant surface-applied N can become positionally unavailable to the SDI corn because of dry surface layers and insufficient root exploration (Lamm, 2014).

Summary of SDI for corn

As corn is by far the major irrigated crop in the USA, it has garnered considerable SDI research attention even though it is considered to be a lower-value commodity crop. Subsurface drip irrigation of corn has generally not increased corn grain yields over well-managed alternative irrigation systems, but may stabilize yields at a greater level when corn is deficit irrigated. Dripline depth for corn often is approximately 0.30 m and dripline spacing is usually an alternate row middle arrangement. Both of these aspects are important in obtaining cost-effective, multiple-year systems that are necessary for SDI adoption for lower-value commodity crops such as corn. Economic competitiveness for SDI as compared to center pivot sprinklers arises from SDI being able to irrigate a greater fraction of the field area and is improved when corn selling prices and grain yields are greater. As is the case with several other crops, SDI offers opportunities for improved conjunctive management of nutrients and irrigation.

Table 3. Corn grain yield, Mg/ha as affected by irrigation method and average yield increase attributable to SDI.

Research report	Additional factors or comments ¹	Soil type	SDI dripline depth (m)	Subsurface Drip (SDI)	Surface drip (DI)	Surface gravity	Sprinkler	Average increased yield using SDI over all alternative methods, %
Abou Kheira, 2009	100% ET	Loamy	0.25	7.3	7.4	11.6	-	-23
	80% ET			4.1	7.2	9.5	-	-51
	60% ET			3.4	4.5	8.0	-	-45
Abuarab et al., 2013	SDI without air injection	Sandy clay loam	0.20	11.3	9.2	-	-	23
	SDI with air injection	loam		12.7	9.2	-	-	38
Adamsen, 1992	Nonsodic water	Loamy sand	0.41	9.9	-	-	10.1	-2
	Sodic water	sand		10.0	-	-	9.9	1
Camp, et al., 1989	0.76 m dripline spacing	Loamy sand	0.30	11.7	12.1			-3
Douh and Boujelben, 2011	SDI depth, 0.05 m	Sandy clay	See treatment depth	10.1	10.4	-	-	-3
	SDI depth, 0.20 m			12.4	10.4	-	-	19
	SDI depth, 0.35 m			13.5	10.4	-	-	30
Hassanli, et al., 2009	Fresh water	Clay loam	0.15 - 0.20	11.8	11.4	9.6	-	9
	Effluent			12.0	11.6	10.4	-	12
Howell, et al., 1997	Daily, 100% ET	Clay loam	0.30	12.9	12.8	-	-	1
	Weekly, 100% ET			13.1	13.0	-	-	0
	Daily, 67% ET			11.4	11.6	-	-	-2
	Weekly, 67% ET			11.6	10.9	-	-	7
	Daily, 33% ET			6.5	6.4	-	-	1
	Weekly, 33% ET			6.6	7.0	-	-	-6
Lamm, 2004	All years, irrigation capacity, 6.4 mm/d	Silt loam	0.40 - 0.45	15.5	-	-	15.4	0
	All years, irrigation. capacity, 4.2 mm/d			14.7	-	-	14.8	-1
	All years, irrigation. capacity, 3.2 mm/d			13.7			14.0	-3
	Normal years, irrigation. capacity, 6.4 mm/d			17.1	-	-	15.7	8
	Normal years, irrigation capacity, 4.2 mm/d			16.5	-	-	15.5	6
	Normal years, irrigation capacity, 3.2 mm/d			16.0			15.6	2
	Drought years, irrigation capacity, 6.4 mm/d			14.3	-	-	15.2	-6
	Drought years, irrigation. capacity, 4.2 mm/d			13.3	-	-	14.3	-7
	Drought years, irrigation. capacity, 3.2 mm/d			11.9			11.9	0
Mitchell and Sparks, 1982	Chiseled treatments	Loamy sand	0.36	10.6	9.1	-	-	16
Oron et al., 1991a	One dripline/bed	Silt loam	0.30	10.6	10.2	-	-	4
	Two driplines/bed	loam		10.1	10.0	-	-	1
Schultz, 2000	Single SDI dripline shown here	Clay loam	0.15 to 0.20	9.3	9.5	8.3	-	4
Stone et al., 2008	1.0 m dripline spacing, DI and SDI	Loamy sand	0.3	5.7	5.6			2
	2.0 m dripline spacing, DI and SDI			5.1	5.2			-2
Average SDI increase across all studies and treatments								3
Average SDI increase over surface gravity irrigation across all studies								-7
Average SDI increase over all sprinkler irrigation treatments across all studies								0
Average SDI increase over DI across all studies								4

Onion

Onion (*Allium cepa* L.) is grown with shallow (< 0.15 m) SDI systems in several areas of the United States, particularly in the Pacific Northwest states of Oregon and Washington. In many respects these systems are very similar to DI production and often the SDI systems are replaced annually for this higher value crop. The current commercial use of SDI for onions is focused primarily on increasing the fraction of larger onions that can command a premium market price and has a much smaller focus on water conservation (Shock, et al., 2013a, Peters, 2015).

System comparisons for onion

Only four studies were found that in the literature that allowed equal or nearly equal comparisons of onion yield under SDI and alternative irrigation systems. Increased soil water availability and reduced soil strength on soils wetted by SDI were contributing factors in higher onion yields on a fine sandy loam soil in India (Abrol and Dixit, 1972). They found SDI onion yields were 24% greater, average diameter was 7% greater and irrigation was 7% less than the best check basin surface irrigation method. When compared to check basin surface irrigation method with the same total irrigation amount, SDI onion yield and diameter were 70 and 25% greater, respectively. In a study in California on a silty loam soil, no statistical differences in onion yield/water use relationships for SDI and DI were reported (Hanson and May, 2003b). At the highest irrigation level (120% of baseline amount) DI numerically had 7% greater total yield, 13% less colossal and 16% greater jumbo yield than SDI. In comparison for the driest irrigation treatment, SDI numerically had 10% greater total yield, 316% greater colossal yield, and 6% less jumbo yield. SDI produced 48% more colossal, 28% more jumbo, and 32% less medium-sized onions than furrow irrigation in a two-year study on a clay loam soil in Colorado (Halvorson, et al., 2008). They concluded that SDI could improve economic returns, and irrigation and nitrogen use efficiencies of Colorado onion production systems. On a silt loam soil in eastern Oregon, Feibert et al. (1995) found SDI onion yield to be 5% and 19% greater for sprinkler and furrow irrigation, respectively. Since that time, shallow (approximately 0.10 m depth) SDI for onion has steadily grown in eastern Oregon and was anticipated to reach 50% of the irrigated onion area by 2013 (Shock et al., 2013a). Extensive research on SDI onion in this region along with grower guidelines have been summarized by Shock et al., (2013a and b).

Dripline depth and irrigation management for onion

Onion is a shallow-rooted crop and adequate soil water must be available near the base plate of the bulb for vigorous growth and development. Thus, dripline depth and irrigation onset and frequency criteria must be carefully considered for SDI onions. Dripline depths of 0, 0.05, 0.10, 0.15, 0.20, and 0.30 m were evaluated for onion production on a sandy loam soil in India (Patel and Rajput, 2009). They found maximum yield and maximum irrigation water use efficiency for a dripline depth of 0.10 m across three irrigation levels (60, 80 or 100% ET) although in most cases the 0.05 and 0.15 m depths were statistically similar. Many SDI onion reports indicated dripline depths of 0.05 to 0.10 m (Feibert et al., 1995; Shock et al., 2005; Halvorson et al., 2008; Shock, et al., 2013a and b; and Pinto et al., 2014), while a few indicate depths of 0.13 to 0.15 m (Shock et al., 2000; Shock et al., 2004, and Enciso et al., 2009), and a few indicate depths as great as 0.18 to 0.23 m (Abrol and Dixit, 1972; Bucks et al., 1981; Al-Jamal et al., 2001; Hanson et al., 2003, and Hanson and May, 2003b). It appears from these studies that a dripline depth of approximately 0.10 m is most prevalent. Deeper dripline depths would be advantageous in the harvesting operation and to increase the possibility of multiple-year usage, but deeper dripline depths can decrease irrigation efficiency through deep percolation losses (Al-Jamal et al., 2001) and may not adequately wet the entire onion crop bed on some soil types (Hanson and May, 2003b). Soil water potentials (SWP) of -10, -20, -30, -50 and -70 kPa measured beneath the onion row at the 0.2 m depth as controlled by an automated high frequency irrigation system were evaluated as SDI management criteria on a silt loam soil in Oregon (Shock et al., 2000). They found that total and colossal yield increased with increasing SWP in both years of the study, but that marketable yield tended to be best in the calculated SWP range of -17 to -21 kPa due to greater onion decomposition in storage. In this report and in later grower recommendations (Shock et al., 2013 b) they conclude a SWP management criteria of approximately -20 kPa is most appropriate for onion production in eastern Oregon on silt loam soils. In a later study (Shock et al., 2005) at the same location, a target SWP of -20 kPa was compared with varying fixed irrigation event amounts (1.6, 3.2, 6.4 and 13 mm) resulting in varying irrigation frequencies (i.e., 564, 269, 121 and 60 events for the 107 day season, respectively). They concluded that irrigation event amounts of 6.4 or 13 mm were superior to the smaller amounts in total, super colossal, and colossal onion yield and that greater irrigation intensity would result in greater leaching losses. Marketable onion yields were 15% greater for daily as compared to weekly irrigation in an Arizona study (Bucks et al., 1981) for a 0.15 m dripline depth. They also reported they were able to successfully harvest the onion without damaging the SDI system which would allow for multiple-season usage. Irrigation scheduled by ET levels (50, 75, and 100%) were compared to SWP levels (-50, -30 and -20 kPa) for SDI onion production in the Lower Rio Grande region of Texas on a sandy clay loam soil (Enciso et al, 2009). They found that SWP levels of -20 and -30 kPa and ET levels of 75 and 100% all resulted in statistically similar yields. Numerically, slight increases in total, colossal, and jumbo yield were reported for the -20 kPa SWP and 100% ET levels although these treatments required 12% more irrigation than the intermediate irrigation levels. Irrigation frequencies of 2 events/week, 1 event/day, or 2 events/day were statistically

similar in SDI onion yield and were significantly greater than the yield for weekly irrigation on a sandy loam soil in California (Hanson et al., 2003). Summarizing these overall onion dripline depth and irrigation management results, it appears that dripline depth for onion generally is in the range of 0.1 to 0.15 m, and that frequent water applications are necessary (daily or at least twice weekly) are necessary to keep SWP near -20 kPa in the vicinity of the crop row.

Summary of SDI for onion

Subsurface drip irrigation for onion generally provides modest yield improvements over alternative irrigation systems, but particularly improves onion quality through increases in the fraction of larger onions. Shallow SDI systems are necessary to ensure adequate soil water in the vicinity of the bulb and the inherent ability of SDI to precisely manage irrigation inputs is advantageous.

Conclusions

The rationale for using SDI for cotton, processing tomato, corn, and onion production varies. Research results indicate moderate or larger yield increases over alternative irrigation systems for cotton, processing tomato and onion with the latter two obtaining differences particularly in marketable yield and quality. Yield increases for SDI field corn on average were small or nonexistent compared to other irrigation systems. The rationale for SDI field corn is more related to stabilizing grain yields when the crop is deficit irrigated and also having less nonirrigated land area for quadrilateral fields than center pivot sprinkler irrigation. Research indicated that precise water and/or nutrient management was a benefit of SDI for all four crops. Frequency of SDI (daily to weekly) was not important for the deeper-rooted commodity crops (cotton and corn) but tended to be more beneficial for the vegetable crops tomato and onion. Dripline spacing using an alternate row middle placement typically was acceptable for cotton and corn. Dripline depth for cotton and corn was approximately 0.3 m and was generally shallower for the vegetable crops ranging from approximately 0.15 to 0.35 m and 0.10 to 0.15 m for tomato and onion, respectively. The use of subsurface irrigation in the USA continues to increase and currently is concentrated in ten states.

Acknowledgements

Contribution number 15-304-J from the Kansas Agricultural Experiment Station.

References

- Abou Kheira, A. A. 2009. Comparison among different irrigation systems for deficit-irrigated corn in the Nile Valley. *Agric. Eng. Int'l: CIGR Ejournal*. Manuscript LW 08 010. Vol. XI. Febr. 2009. 25 pp.
- Abrol, I. P. & Dixit, S. P. 1972. Studies of the drip method of irrigation. *Expl. Agric.* 8:172-175.
- Abuarab, M., Mostafa, E., & Ibrahim, M. 2013. Effect of air injection under subsurface drip irrigation on yield and water use efficiency of corn in a sandy clay loam soil. *J. Adv. Res.* (2013)4:493-499. Elsevier B.V. <http://dx.doi.org/10.1016/j.jare.2012.08.009>.
- Adamsen, F. J. 1992. Irrigation method and water quality effects on corn yield in the mid-Atlantic coastal plain. *Agron J.* 84:837-843.
- Al-Jamal, M. S., Ball, S., & Sammis, T. W. 2001. Comparison of sprinkler, trickle and furrow irrigation efficiencies for onion production. *Agric. Water Mgt.* 46(2001):253-266.
- Aloni, N. I., Eizencot, A., Steinberg, Y., Levy, U., Yotal, Y., & Golan, Z. 2000. Fertigation management and salinity monitoring of cotton cultivated in deep subsurface drip irrigation system. In: *Proc. Nat'l. Conf. and Exhibition of Irrigation Assoc. Australia, May 23-25, 2000, Melbourne, Australia.* pp. 153-157.
- ASAE S526.3. 2007. ASAE Standard S526.3, SEP07, Soil and Water Terminology. ASAE, St. Joseph, Michigan. 55 pp.
- Ayars, J. E., Phene, C. J., Hutmacher, R. B., Davis, K. R., Schoneman, R. A., Vail, S. S., & Mead, R. M. 1999. Subsurface drip irrigation of row crops: a review of 15 years of research at the Water Management Research Laboratory. *Agric. Water Mgt.* 42:1-27.
- Bar-Yosef, B., Sagiv, B., & Markovitch, T. 1989. Sweet corn response to surface and subsurface trickle phosphorous fertigation. *Agron. J.* 81:443-447.
- Batchelor, C. H., Lovell, C. J., Murata, M., & McGrath, S. P. 1994. Improving water use effectiveness with subsurface irrigation. *Aspects Appl. Biol.* 38:269-278.
- Bauer, P. J., Hunt, P. G., & Camp, C. R. 1997. In-season evaluation of subsurface drip and nitrogen-application method for supplying nitrogen and water to cotton. *J. Cotton Sci.* 1:29-37.
- Ben-Asher, J. & Silberbush, M. 1992. Root distribution under trickle irrigation: Factors affecting distribution and comparison among methods of determination. *J. Plant Nutrition* 15(6 & 7):783-794.
- Ben-Gal, A. & Dudley, L. M. 2003. Phosphorus availability under continuous point source irrigation. *Soil Sci. Soc. Amer. J.* 67:1449-1456.
- Bhattarai, S. P., Huber, S., & Midmore, D. J. 2004. Aerated subsurface irrigation gives growth and yield benefits to zucchini, vegetable soybean and cotton in heavy clay soils. *Ann. Appl. Biol.* 144:285-298.
- Bhattarai, S. P., McHugh, A. D., Lotz, G. & Midmore, D. J. 2005. The response of cotton to subsurface drip and furrow irrigation in a vertisol. *Expl. Agric.* 42(1):29-49.
- Bogle, C. R., Hartz, T. K. & Nunez, C. 1989. Comparison of subsurface trickle and furrow irrigation on plastic-mulched and bare soil for tomato production. *J. Amer. Hort. Sci.* 114(1):40-43.

- Bordovsky, J. P. & Lyle, W. M. 1998. Cotton irrigation with LEPA and subsurface drip systems on the southern high plains. In Proc. Beltwide Cotton Conf., San Diego, CA., Jan. 5-9, 1998. pp. 409-412.
- Bordovsky, J. P. & Porter, D. 2003. Cotton response to pre-plant irrigation level and irrigation capacity using spray, LEPA and subsurface drip irrigation. Presented at the annual int'l. mtg. of the ASAE, Las Vegas, Nevada, July 27-30, 2003. ASAE Paper No. 032008. 11 pp.
- Bordovsky, J. P., Emerson, C. L. W., & Mustian, J. T. 2012a. Irrigation interval effects on cotton production using subsurface drip systems. Presented at the 2012 ASABE Annual International Meeting. ASABE Paper No. 12-1337401. ASABE, St. Joseph, MI. 10 pp.
- Bordovsky, J.P., Cranmer, A. M., Colaizzi, P. D., Lamm, F. R., Evett, S. R. & Howell, T. A. 2012b. Investigating strategies to improve crop germination when using SDI. In: Proc. 24th annual Central Plains Irrigation Conference Feb. 21-22, 2012, Colby, Kansas. Available from CPIA, 760 N. Thompson, Colby, Kansas. pp. 112-127.
- Bosch, D. J., Powell, N. L., & Wright, F. S. 1992. An economic comparison of subsurface microirrigation and center pivot sprinkler irrigation. *J. Prod. Agric.* 5(4):431-437.
- Bosch, D. J., Powell, N. L., & Wright, F. S. 1998. Investment returns from three sub-surface microirrigation tubing spacings. *J. Prod. Agric.* 11(3):371-376.
- Bucks, D. A., Erie, L. J., French, O. F., Nakayama, F. S., & Pew, W. D. 1981. Subsurface trickle irrigation management with multiple cropping. *Trans. ASAE* 24(6):1482-1489.
- Caldwell, D. S., Spurgeon, W. E., & Manges, H. L. 1994. Frequency of irrigation for subsurface drip-irrigated corn. *Trans. ASAE* 37(4):1099-1103.
- Camp, C. R. 1998. Subsurface drip irrigation: A review. *Trans. ASAE* 41(5):1353-1367.
- Camp, C. R., Bauer, P. J., & Hunt, P. G. 1997. Subsurface drip irrigation lateral spacing and management for cotton in the southeastern Coastal Plain. *Trans. ASAE* 40(4):993-999.
- Camp, C. R., Sadler, E. J., & Busscher, W. J. 1989. Subsurface and alternate-middle micro irrigation for the Southeast Coastal Plain. *Trans. ASAE* 32(2):451-456.
- Camp, C. R., Lamm, F. R., Evans, R. G., & Phene, C. J. 2000. Subsurface drip irrigation: Past, present and future. In: Proc. 4th Decennial National Irrig. Symp., Phoenix, AZ, Nov. 14-16, 2000. pp. 363-372.
- Clark, G. A., Stanley, C. D., & Maynard, D. N. 1993. Surface vs. subsurface drip irrigation of tomatoes on a sandy soil. *Proc. Florida State Hort. Soc.* 106:210-212.
- Colaizzi, P. D., Evett, S. R., & Howell, T. A. 2006. Crop emergence with alternative designs in a Pullman clay loam soil. In: Proc. Southern Conservation Systems Conference, Amarillo, Texas, June 26-28. pp. 16-24.
- Colaizzi, P. D., Evett, S. R., Howell, T. A., & Baumhardt, R. L. 2010. Crop Production Comparison with Spray, LEPA, and Subsurface Drip Irrigation in the Texas High Plains. In: Proc. Fifth Nat'l. Irrigation Symp., Phoenix, Dec. 5-8. ASABE, St Joseph Michigan. 23 pp.
- Constable, G. A. & Hodgson, A. S. 1990. A comparison of drip and furrow irrigated cotton on a cracking clay soil. 3. Yield and quality of four cultivars. *Irrig. Sci.* 11:149-153.
- Constable, G. A., Rochester, I. J., & Hodgson, A. S. 1990. A comparison of drip and furrow irrigated cotton on clay soil. 1. Growth and nitrogen uptake. *Irrig. Sci.* 11:137-142.
- Cote, C. M., K. L. Bristow, P. B. Charlesworth, F. J. Cook, and P. J. Thorburn. 2003. Analysis of soil wetting and solute transport in subsurface trickle irrigation. *Irrig. Sci.* 22:143-156.
- Darusman, Khan, A. H., Stone, L. R., & Lamm, F. R. 1997. Water flux below the root zone vs. drip-line spacing in drip irrigated corn. *Soil Sci. Soc. Am. J.* 61(6):1755-1760. (NOTE to ASABE Darusman is complete name, no initials)
- del Amor, M. A. & del Amor, F. M. 2007. Response of tomato plants to deficit irrigation under surface or subsurface drip irrigation. *J. Appl. Hort.* 97(2):97-100.
- Devasirvatham, V. 2009. A review of subsurface drip irrigation in vegetable production. CRC for Irrigation Futures, Irr. Matters Series No. 03/09. Darling Heights, QLD, Australia. 35 pp.
- Douh, B. & Boujelben, A. 2011. Effects of surface and subsurface drip irrigation on agronomic parameters of maize (*Zea mays* L.) under Tunisian climatic condition. *J. Nat. Prod. Plant Resour.*, 1(3):8-14.
- Dukes, M. D. & Scholberg, J. M. 2005. Soil moisture controlled subsurface drip irrigation on sandy soils. *Appl. Engr. Agric.* 21(1):89-101.
- El-Gindy, A. M. & El-Araby, A. M. 1996. Vegetable crop response to surface and subsurface drip under calcareous soil. In Proc. Int'l Conf. on Evapotranspiration and Irrigation Scheduling, San Antonio, TX, Nov. 3-6, 1996. ASAE, St Joseph Michigan. pp. 1021-1028.
- Enciso, J. M., Colaizzi, P. D., & Multer, W. L. 2005. Economic analysis of subsurface drip irrigation lateral spacing and installation depth for cotton. *Trans. ASAE* 48(1):197-204.
- Enciso, J., Wiedenfeld, B., Jifon, J. & Nelson, S. 2009. Onion yield and quality response to two irrigation scheduling strategies. *Scientia Hort.* 120(2009):301-305.
- Enciso-Medina, J., Unruh, B. L., Colaizzi, P. D. & Multer, W. L. 2003. Cotton response to subsurface drip irrigation frequency under deficit irrigation. *Appl. Engr. Agric.* 19(5):555-558.
- Evett, S. R., Howell, T. A., & Schneider, A.D. 1995. Energy and water balances for surface and subsurface drip irrigated corn. In: Proc. Fifth Int'l. Microirrigation Cong., F. R. Lamm (Ed.), Apr. 2-6, 1995, Orlando Florida. ASAE, St. Joseph, Michigan. pp. 135-140.
- Feibert, E. B. G., Shock, C. C., & Saunders, L. D. 1995. A comparison of sprinkler, subsurface drip, and furrow irrigation of onions. Oregon State Univ. Agr. Expt. Sta. Spec. Rpt. 947:59-67.
- Gilley, J. R. & Allred, E. R. 1974. Infiltration and root extraction from subsurface irrigation laterals. *Trans. ASAE* 17(5):927-933.
- Grabow, G. L., Huffman, R. L., Evans, R. O., Edmisten, K. L., & Jordan, D. L. 2002. Subsurface drip irrigation research on commodity crops in North Carolina. Presented at the Int'l meeting of the ASAE, Chicago, Illinois, July, 28-31. Paper No. 022290. 14 pp.
- Grattan, S. R., Schwankl, L. J., & Lanini, W. T. 1990. Distribution of annual weeds in relation to irrigation method. In: Proc. 3rd National

- Irrigation Symp. Phoenix, AZ, Oct. 28-Nov. 1, 1990. ASAE, St. Joseph, MI. pp. 148-153.
- Halvorson, A. D., Bartolo, M. E., Reule, C. A., & Berrada, A. 2008. Nitrogen effects on onion yield under drip and furrow irrigation. *Agron. J.* 100(4):1062-1069.
- Hanson, B. R. & May, D. M. 2000. Response of processing tomato yield to drip irrigation under saline, shallow groundwater conditions. In: Proc. 4th National Irrigation Symp. Phoenix, AZ, Nov. 14-16, 2000. ASAE, St. Joseph, MI. pp. 385-390.
- Hanson, B. & May, D. M. 2003a. Effect of subsurface drip irrigation on processing tomatoes yield, water table depth and soil salinity. Presented at the annual int'l. mtg. of the ASAE, Las Vegas, Nevada, July 27-30, 2003. ASAE Paper No. 032091. 12 pp.
- Hanson, B. & May, D. M. 2003b. Response of processing and fresh-market onions to drip irrigation. Presented at the annual int'l. mtg. of the ASAE, Las Vegas, Nevada, July 27-30, 2003. ASAE Paper No. 032094. 10 pp.
- Hanson, B. & May, D. M. 2007. The effect of dripline placement on yield and quality of drip-irrigated processing tomatoes. *Irrig. Drainage Syst.* 21:109-118. DOI 10.10007/s10795-007-9023-5.
- Hanson, B.R., May, D. M. & Schwankl, L. J. 2003. Effect of irrigation frequency on subsurface drip irrigated vegetables. *HortTechnology* 13(1):115-120.
- Hanson, E. G., Williams, B. C., Fangmeier, D. D., & Wilke, O. C. 1970. Influence of subsurface irrigation on crop yields and water use. In: Proc. Nat'l Irrigation Symposium, Lincoln, Nebraska, Nov 10-13, 1970. pp. D1-D13.
- Hassanli, A. M., Ebrahimzadeh, M. A., & Beecham, S. 2009. The effects of irrigation methods with effluent and irrigation scheduling on water use efficiency and corn yields in an arid region. *Agri. Water Manage.* 96(2009):93-99. Elsevier ScienceDirect. www.elsevier.com/locate/agwat.
- Hollingsworth, J., Mitchell, J. P., Munk, D.S., Roberts, B. A., & Shresetha, A. 2014. Subsurface drip and overhead irrigation effects on conservation-tilled cotton in the San Joaquin Valley. *J. Crop Improvement* 28:324-344.
- House, E. B. 1918. Irrigation by Means of Underground Porous Pipe. *Colorado Agric. Expt. Sta. Bull.* 240, Fort Collins, Colorado. 42 pp.
- Howell, T. A., Schneider, A. D., & Evett, S. R. 1997. Subsurface and surface microirrigation of corn, Southern High Plains. *Trans ASAE* 40(3):635-641.
- HPWCD. 2014. High Plains Water Conservation District 2013 subsurface drip irrigation locations. Available at http://www.hpwd.org/s/SDI_2013.pdf. Verified, 2-13-15.
- Hulme, P. & O'Brien, B. 2000. Managing subsurface drip irrigation in hardsetting red soil – Implications of small wetted area. In: Proc. Nat'l Conf. and Exhibition of Irrigation Assoc. Australia, May 23-25, 2000, Melbourne, Australia. pp. 385-390.
- Kahlaoui, B., Hachicha, M., Rejeb, S., Rejeb, M. N., Hanchi, B., & Misle, E. 2011. Effects of saline water on tomato under subsurface drip irrigation: Nutritional and foliar aspects. *J. Soil Sci. Plant Nutr.* 11(1):69-86.
- Kamara, L., Zartman, R., & Ramsey, R. H.. 1991. Cotton-root distribution as a function of trickle emitter depth. *Irrig. Sci.* 12:141-144.
- Khalilian, A., Sullivan, M. J., & Smith, W. B. 2000. Lateral depth placement and deep tillage effects in a subsurface drip irrigation system for cotton. In: Proc. 4th Nat'l Irrigation Symposium, Phoenix, Arizona, Nov 14-16, 2000. ASAE. pp 641-646.
- Lamm, F. R. 2004. Comparison of SDI and Simulated LEPA Sprinkler Irrigation for Corn. In Proc. Irrigation Assn. Int'l. Irrigation Technical Conf., November 14-16, 2004, Tampa, FL. Available from Irrigation Assn., Falls Church VA. IA Paper No. IA04-1098. pp 475-485.
- Lamm, F. R. 2014. Irrigation and nitrogen management for subsurface drip irrigated corn - 25 years of K-State's efforts. ASABE paper no. 141914980. Available from ASABE, St. Joseph, MI. 10 pp.
- Lamm, F. R. & Aiken, R. M. 2005. Effect of irrigation frequency for limited subsurface drip irrigation of corn. In: Proc. Irrig. Assoc. Int'l. Tech. Conf., Nov. 6-8, 2005, Phoenix, AZ. Irrigation Assoc., Falls Church, Virginia. IA Paper No. IA05-1264. 10 pp.
- Lamm, F. R. & Camp, C. R. 2007. Subsurface drip irrigation. Chapter 13 in *Microirrigation for Crop Production - Design, Operation and Management*. F. R. Lamm, J. E. Ayars, and F. S. Nakayama (Eds.), Elsevier Publications. pp. 473-551.
- Lamm, F. R. & Trooien, T. P. 2003. Subsurface drip irrigation for corn production: A review of 10 years of research in Kansas. *Irrig. Sci.* 22(3-4):195-200.
- Lamm, F. R. and T. P. Trooien. 2005. Dripline depth effects on corn production when crop establishment is nonlimiting. *Appl. Engr. in Agric.* 21(5):835-840.
- Lamm, F. R., O'Brien, D. M., & Rogers, D. H. 2015. Using the K-State center pivot sprinkler and SDI economic comparison spreadsheet – 2015. In: Proc. 27th annual Central Plains Irrigation Conference, Feb. 17-18, 2015, Colby, Kansas. Available from CPIA, 760 N. Thompson, Colby, Kansas. pp. 108-116.
- Lamm, F. R., Schlegel, A. J. & Clark, G. A. 2004. Development of a best management practice for nitrogen fertigation of corn using SDI. *Appl. Engr in Agric.* 20(2):211-220.
- Lamm, F. R., Trooien, T. P., Manges, H. L., & Sunderman, H. D. 2001. Nitrogen fertilization for subsurface drip-irrigated corn. *Trans. ASAE* 44(3):533-542.
- Lamm, F. R., Stone, L. R., Manges, H. L., & O'Brien, D. M. 1997. Optimum lateral spacing for subsurface drip-irrigated corn. *Trans. ASAE* 40(4):1021-1027.
- Lamm, F. R., Colaizzi, P. D., Bordovsky, J. P., Trooien, T. P., Enciso-Medina, J., Porter, D. O., Rogers, D. H., & O'Brien, D. M. 2010. Can Subsurface Drip Irrigation (SDI) be a Competitive Irrigation System in the Great Plains Region for Commodity Crops? In: Proc. 5th National Decennial Irrigation Conf., ASABE and the Irrigation Association, Phoenix, Arizona, December 5-8. 15 pp.
- Lamm, F. R., Bordovsky, J. P., Schwankl, L. J., Grabow, G. L., Enciso-Medina, J., Peters, R. T., Colaizzi, P. D., Trooien, T. P., & Porter, D. O. 2012. Subsurface drip irrigation: Status of the technology in 2010. *Trans. ASABE* 55(2): 483-491.
- Machado, R. M. A., do Rosario, M., & Oliveira, G. 2005. Tomato root distribution, yield and fruit quality under different subsurface drip irrigation regimes and depths. *Irrig. Sci.* 24:15-24.
- Machado, R. M. A., do Rosario, M., Oliveira, G., & Portas, C. A. M. 2003. Tomato root distribution, yield and fruit quality under subsurface

- drip irrigation. *Plant and Soil* 255:333-341.
- Marouelli, W. A. & W. L. C. Silva, W. L. C. 2002. Profundidade de instalacao da linha de gotejadores em tomateiro para processamento industrial. *Horticultura Brasileira*, Brasilia 20(2):206-210.
- Martinez-Hernandez, J. J., Bar-Yosef, B., & Kafkafi, U. 1991. Effect of surface and subsurface drip fertigation on sweet corn rooting, uptake, dry matter production and yield. *Irrig. Sci.* 12:153-159.
- Mitchell, W. H. 1981. Subsurface irrigation and fertilization for corn. *Agron. J.* 73(6):913-916.
- Mitchell, W. H. & Sparks, D. L. 1982. Influence of subsurface irrigation and organic additions on top and root growth of field corn. *Agron. J.* 74(6):1084-1088.
- Mitchell, W. H. & Tilmon, H. D. 1982. Underground trickle irrigation – The best system for small farms? *Crops and Soils* 34(5):9-13.
- Mitchell, W. H., McVaine, J. E., & Mueller, J. P. 1969. Subsurface irrigation with perforated plastic tubing. *Agric. Ext. Service Bull.* No. 99. 23 pp.
- Montazar, A. 2015. Personal communication from agricultural water management specialist, Dept. of Plant Sciences, Univ. California-Davis. January 7, 2015.
- Montemayor Trejo, J. A., Gómez Monsivais, A. O., Olague Ramírez, J., Zermeño González, A., Ruiz Cerda, E., Fortis Hernández, M., Salazar Sosa, E., & Aldaco Nuncio, R. 2006. Effect of three driptape installation depths on water use efficiency and yield parameters in forage maize. *Téc. Pecu. Méx.*44(3):359-364.
- Najafi, P. & Tabatabaei, S. H. 2007. Effect of using subsurface drip irrigation and ET-HS model to increase WUE in irrigation of some crops. *Irrig. and Drain.* 56:477-486.
- Nuti, R.C., Casteel, S. N., Viator, R. P., Lanier, J. E., Edmisten, K. L., Jordan, D. L., Grabow, G. L., Barnes, J. S., Mathews, J. W., & Wells, R. 2006. Management of cotton grown under overhead sprinkle and sub-surface drip irrigation. *J. Cotton Science* 10:76-88.
- O'Brien, D. M., Rogers, D. H., Lamm, F. R., & Clark, G. A. 1998. An economic comparison of subsurface drip and center pivot sprinkler irrigation systems. *Appl. Engr. Agric.* 14(4):391-398.
- Oron, G., DeMalach, Y., Hoffman, Z., & Cibotaru, R. 1991a. Subsurface microirrigation with effluent. *ASCE J. Irrig Drain. Engr.* 117(1):25-36.
- Oron, G., DeMalach, Y., Hoffman, Z., Keren, Y., Hartman, H., & Plazner, N. 1991b. Wastewater disposal by sub-surface trickle irrigation. *Water Sci. Tech.* 23:2149-2158.
- Peters, R. T. 2015. Personal communication from agricultural water management specialist, Washington State Univ. January 5, 2015.
- Patel, N. & Rajput, T. B. S. 2009. Effect of subsurface drip irrigation on onion yield. *Irrig. Sci.* 27:97-108. DOI 10.1007/s000271-008-0125-0.
- Pablo, R. G., O'Neill, M. K., McCastin, B. D., Remmenga, M. D., Keenan, J. G., & Onken, B. M. 2007. Evaluation of corn grain yield and water use efficiency using subsurface drip irrigation. *J. Sust. Agric.* 30(1):153-172. DOI: 10.1300/J064v30n01_10.
- Pinto, J. M., Shock, C. C., & Feibert, E. B. G. 2014a. Optimal practices for drip irrigated onion differ based on growers' market opportunities. In: *Proc. 2014 Irrigation Association Tech. Conf.*, Phoenix, AZ, November 19-20, 2014. Available from the IA, Falls Church, VA. 5 pp.
- Phene, C. J., DeTar, W. R. & Clark, D. A. 1992. Real-time irrigation scheduling of cotton with an automated pan evaporation system. *Appl. Engr. Agric.* 8(6):787-793.
- Phene, C.J., Huttmacher, R. B., Davis, K. R., & McCormick, R. L. 1990. Water-fertilizer management of processing tomatoes. *Acta Hort.* 277:137-143.
- Phene, C. J., Davis, K. R., Huttmacher, R. B. & McCormick, R. L. 1987. Advantages of subsurface irrigation for processing tomatoes. *Acta Hort.* 200:101-114.
- Phene, C. J., Davis, K. R., Huttmacher, R. B., Bar-Yosef, B., Meek, D. W., & Misaki, J.. 1991. Effect of high frequency surface and subsurface drip irrigation on root distribution of sweet corn. *Irrig. Sci.* 12:135-140.
- Philip, J. R. 1991. Effects of root and subirrigation depth on evaporation and percolation losses. *Soil Sci. Soc. Am. J.* 55:1520-1523.
- Plaut, Z., Carmi, A., & Grava, A. 1996. Cotton root and shoot responses to subsurface drip irrigation and partial wetting of the upper soil profile. *Irrig. Sci.* 16:107-113.
- Plaut, Z., Rom, M., & Meiri, A. 1985. Cotton response to subsurface trickle irrigation. In: *Proc. 3rd Int'l Drip Trickle Irrigation Congress*, Fresno, California, Nov 18-21. ASAE, St. Joseph MI. Vol. 2, 887-895.
- Powell, N. L. & Wright, F. S. 1993. Grain yield of subsurface microirrigated corn as affected by irrigation line spacing. *Agron. J.* 85:1165-1170.
- Rodríguez-Sinobas, L. & Gil-Rodríguez, M. 2012. A review of subsurface drip irrigation and its management. In: *Water Quality, Soil and Managing Irrigation of Crops*, T. S.hui Lee (Ed.), ISBN: 978-953-51-0426-1, InTech. pp. 171-194. Available from: <http://www.intechopen.com/books/water-quality-soil-and-managingirrigation-of-crops/a-review-of-subsurface-drip-irrigation-and-its-management> . Verified, 2-13-15.
- Schwankl, L. J., Grattan, S. R., & Miyao, E. M. 1990. Drip irrigation burial depth and seed planting depth effects on tomato germination. In: *Proc. 3rd Nat'l Irrigation Symp.* Phoenix, AZ, October 28-November 1, 1990. ASAE, St. Joseph, Michigan. pp. 682-687.
- Schulz, M. 2000. Microirrigation of maize. In: *Proc. Nat'l Conf. and Exhibition of Irrigation Assoc.* Australia, May 23-25, 2000, Melbourne, Australia. pp. 398-405.
- Shock, C. C., Feibert, E. B. G., & Saunders. L. D. 2000. Irrigation criteria for drip-irrigated onions. *HortScience* 35(1):63-66.
- Shock, C. C., Feibert, E. B. G., & Saunders. L. D. 2004. Plant population and nitrogen fertilization for subsurface drip-irrigated onion. *HortScience* 39:1722-1727.
- Shock, C. C., Feibert, E. B. G., & Saunders. L. D. 2005. Onion response to drip irrigation intensity and emitter flow rate. *HortTechnology* 15:652-659.
- Shock, C. C., Feibert, E. B. G., & Pinto, J. M. 2013a. Review of two decades of progress in the development of successful drip irrigation for

- onions. In: Proc. 2013 Irrigation Association Tech. Conf., Austin, Texas, November 5-8, 2013. Available from the IA, Falls Church, VA. CD-Rom. 20 pp.
- Shock, C. C., Flock, R., Feibert, E. Shock, C. A., & Klauzer, J. 2013b. Drip irrigation guide for onion growers. Oregon St. Univ. Ext. publication, EM 8901. 8 pp.
- Shrestha, A., Mitchell, J. P., & Lanini, W. T. 2007. Subsurface drip irrigation as a weed management tool for conventional and conservation tillage tomato (*Lycopersicon esculentum* Mill.) production in semi-arid agroecosystems. *J. Sustainable Agric.* 31(2):91-112.
- Smith, R. B., Oster, J. D., & Phene, C. J. 1991. Subsurface drip produced highest net return in Westlands area study. *California Agric.* 45(2):8-11.
- Sorensen, R. B., Bader, M. J., & Wilson, E. H. 2004. Cotton yield and grade response to nitrogen applied daily through a subsurface drip irrigation system. *Appl. Engr. in Agric.* 20(1):13-16.
- Sorensen, R. B., Butts, C. L., & Lamb, M. C. 2013. Corn yield response to deep subsurface drip irrigation in the southeast. Online. *Crop Management.* 9 pp. Doi:10.1094/CM-2013-0122-01-RS
- Sorensen, R. B., Butts, C. L., & Nuti, R. C. 2011. Deep subsurface drip irrigation for cotton in the Southeast. *J. Cotton Science* 15:233-242.
- Spurgeon, W. E., Manges, H. L., & Makens, T. P. 1991. Drip line spacing and plant population for corn. Presented at the annual int'l. winter mtg. of the ASAE, Chicago, Illinois, December 17-20, 1991. ASAE Paper No. 912592. 12 pp.
- Stone, K. C., Bauer, P. J., Busscher, W. J., & Millen, J. A. 2008. Narrow row corn production with subsurface drip irrigation. *Appl. Engr. Agric.* 24(4):455-464.
- Tan, C. S., Zhang, T. Q., Reynolds, W. D., Drury, C. F., & Liptay, A. 2003. Farm-scale processing tomato production using surface and subsurface drip irrigation and fertigation. Presented at the annual int'l. mtg. of the ASAE, Las Vegas, Nevada, July 27-30, 2003. ASAE Paper No. 032092. 13 pp.
- Tarkalson, D. D. & Payero, J. O. 2008. Comparison of nitrogen fertilization methods and rates for subsurface drip irrigated corn in the semi-arid Great Plains. *Trans. ASABE* 51(5):1633-1643.
- Taylor, R., Parker, D., & Zilberman, D. 2014. Contribution of University of California Cooperative Extension to drip irrigation. *Univ. of California Giannini Foundation of Agric. Econ., ARE Update* 18(2):58.
- Thomas, A. W., Kruse, E. G., & Duke, H. R. 1974. Steady infiltration from line sources buried in the soil. *Trans. ASAE* 17(1):125-133.
- Thompson, W. J. & Enciso, J. M. 2002. Why subsurface drip irrigation (SDI)? An economic analysis. In: Proc. Beltwide Cotton Conference, Atlanta, Georgia, January 8-12. 3 pp.
- Thorburn, P. J., Cook, F. J., & Bristow, K. L. 2003. Soil-dependent wetting from trickle emitters: Implications for system design and management. *Irrig. Sci.* 22:121-127.
- Tollefson, S. 1985. Subsurface drip irrigation of cotton and small grains. In: Proc. 3rd Int'l Drip Trickle Irrigation Congress, Fresno, California, Nov 18-21. ASAE, St. Joseph MI. Vol. 2, 887-895.
- USDA-NASS. 2004. Farm and ranch irrigation survey (2003). In: 2002 Census of Agriculture, Vol. 3, Special Studies, Part 1 (AC-02-SS-1). 216 pp.
- USDA-NASS. 2010. Farm and ranch irrigation survey (2008). In: 2007 Census of Agriculture, Vol. 3, Special Studies, Part 1 (AC-07-SS-1). 268 pp.
- USDA-NASS. 2014. Farm and ranch irrigation survey (2013). In: 2012 Census of Agriculture, Vol. 3, Special Studies, Part 1 (AC-12-SS-1). 266 pp.
- Whitaker, J. R., Ritchie, G. L., Bednarz, C. W., & Mills, C. I. 2008. Cotton subsurface drip and overhead irrigation efficiency, maturity, yield, and quality. *Agron. J.*, 100:1763-1768.
- Zetzsche, J. B. & Newman, J. S. 1966. Subirrigation with plastic pipe. *Agric. Engineering.* 47(1):74-75.