

You Can Reduce Your Risk of Colon Disease



- Don't smoke
 - Diet and Physical Activity Recommendations:
 - Eat three or more cups of a variety of fruits and vegetables
 - At least half of your grain foods should be whole grain foods
 - Eat 3 cups of cooked dried beans each week
 - Eat seafood frequently
 - Limit the amount of added sugars in your diet
 - Eat only moderate amounts of red meat and meats that are smoked, cured or high in fat or sodium
 - Be physically active 30 or more minutes on at least five days each week
 - If overweight, lose weight
 - Avoid drinking excessive amounts of alcohol
-

You Can Reduce Your Risk of Colon Disease



- Don't smoke
- Diet and Physical Activity Recommendations:
 - Eat three or more cups of a variety of fruits and vegetables
 - At least half of your grain foods should be whole grain foods
 - Eat 3 cups of cooked dried beans each week
 - Eat seafood frequently
 - Limit the amount of added sugars in your diet
 - Eat only moderate amounts of red meat and meats that are smoked, cured or high in fat or sodium
 - Be physically active 30 or more minutes on at least five days each week
 - If overweight, lose weight
 - Avoid drinking excessive amounts of alcohol

You Can Reduce Your Risk of Colon Disease

- Take a daily multivitamin and mineral supplement with folate
- Take a supplement with vitamin D and calcium

About one in three deaths from colon and rectal cancer could be avoided if men regularly got a screening test:

Screening Tests Recommended for those ages 50 years and older

- Fecal occult blood test (annually)
- Digital rectal exam (annually)
- Sigmoidoscopy (every 5 years)
- Double contrast barium enema (every 5-10 years)
- Colonoscopy (every 10 years)

Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, credit Mary L. Meck Higgins and Kimberly Shafer, *Men's Health*, 2006, Department of Human Nutrition, **Kansas State University Agricultural Experiment Station and Cooperative Extension Service, Manhattan, Kansas**. K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Fred A. Cholick, Director.

You Can Reduce Your Risk of Colon Disease

- Take a daily multivitamin and mineral supplement with folate
- Take a supplement with vitamin D and calcium

About one in three deaths from colon and rectal cancer could be avoided if men regularly got a screening test:

Screening Tests Recommended for those ages 50 years and older

- Fecal occult blood test (annually)
- Digital rectal exam (annually)
- Sigmoidoscopy (every 5 years)
- Double contrast barium enema (every 5-10 years)
- Colonoscopy (every 10 years)

Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, credit Mary L. Meck Higgins and Kimberly Shafer, *Men's Health*, 2006, Department of Human Nutrition, **Kansas State University Agricultural Experiment Station and Cooperative Extension Service, Manhattan, Kansas**. K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Fred A. Cholick, Director.